

The use of propolis extract for the storage of quail eggs

G. C. Akpınar,^{*,1} S. Canogullari,[†] M. Baylan,[‡] S. Alasahan,[§] and A. Aygun[#]

^{*}Department of Animal Science, Agriculture Faculty, Mustafa Kemal University, Hatay, Turkey, 31034; [†]Animal Production and Technologies, Agricultural Science and Technologies Faculty, Nigde University, Nigde, Turkey 51240; [‡]Department of Animal Science, Agriculture Faculty, Cukurova University, Adana, Turkey, 01330; [§]Department of Animal Sciences, Veterinary Faculty; Mustafa Kemal University, Hatay, Turkey, 31034; and [#]Department of Animal Science, Agriculture Faculty, Selcuk University, Konya, Turkey, 42075.

Primary Audience: Poultry Researchers

SUMMARY

This study was carried out to evaluate the use of propolis extract on egg storage time of table quail (*Coturnix coturnix Japonica*) eggs. Propolis is a sticky, resinous, dark-colored material that honey bees collect from young plants. In the experiment, 600 quail eggs were used and eggs were coated with various concentrations of propolis extract (0%: group I, 5%: group II, 10%: group III, 15%: group IV) and treated with 70% ethyl alcohol (group V). A 5-week storage period was implemented, and 120 eggs were used for each group. Consequently, the effects of storage time and shell treatments on storage time and the effects of treatments on the interior quality of eggs were determined. The results of the study confirmed highly significant differences between weekly changes in egg weight loss, albumen-yellow indexes, and Haugh units ($P < 0.001$). While the difference in yolk index between groups was insignificant, the difference between groups with respect to albumen index ($P < 0.001$), Haugh units ($P < 0.001$) albumen pH ($P < 0.001$), and egg weight loss ($P < 0.001$) were significant. The significance of the overall difference with regard to Haugh units varied among groups; for example, group IV showed the highest value of 87.73%, followed by group III (87.69%), group II (86.97%), group V (85.53%), and group I (85.21%). Albumen pH levels were increased with increasing storage time for each treatment group. The best egg protection results in terms of interior quality were obtained in eggs coated with 10% and 15% propolis extract during storage.

Key words: egg quality, propolis, quail egg, storage time.

2015 J. Appl. Poult. Res. 00:1–9
<http://dx.doi.org/10.3382/japr/pfv043>

DESCRIPTION OF PROBLEM

Eggs are naturally protected against microbial spoilage to some extent thanks to their biochemical structure and composition. However,

significant quality losses may occur depending on the interval between egg collection time and consumption time if necessary measures are not taken [1].

Albumen height and albumen pH are important characteristics in the determination of the internal quality of commercial eggs. Of these

¹Corresponding author: gulsenankara@gmail.com

properties, albumen height decreases, whereas albumen pH value increases with an increase in storage time [2–4]. The albumen pH value is between 7.6 and 7.9 following oviposition and may rise to 9.7 during storage time [5–9]. This happens because of the increased carbon dioxide loss from albumen through the shell [1,5–8].

Measures taken to prolong the egg storage time are usually based on a limitation of gas permeability due to coating of the pores at the shell surface [8,10]. Coating the egg shell protects the natural structure of the egg by reducing CO₂ and water loss from the albumen. For this reason some conservation methods, including oil coating, storing at low temperatures, freezing, storing, and drying, have been used for the prolonged protection of shell life, along with other methods [11].

In addition, some natural and risk-free products in terms of human health have been used in the coating of egg shells, such as chitosan, whey protein, shellac [12], and propolis [9].

Propolis is a sticky, gummy, resinous substance with a strange odor, which is collected by worker honeybees (*Apis mellifera*) from the young shoots and buds of certain trees and shrubs [13,14]. Propolis has immunostimulating, antibacterial, antioxidant, antimicrobial, and antiparasitic properties [15–21], and it is normally used by honeybees (*Apis mellifera*) to repair the slits and cracks inside hives and to embalm dead honeybees or other dead bugs that could not be removed from hives. Additionally, use of propolis during the brood period prevents water loss by evaporation through cracks and crevices [13,14].

Propolis is also used for the protection of various agricultural products during storage because of its antimicrobial and embalming properties. Torre et al. [22] and Ozdemir et al. [23], covered strawberry and mandarin with propolis to inhibit *Botrytis cinerae* Pers development and to prevent weight loss during storage. Owing to its antimicrobial effects, propolis has also been used to disinfect quail hatching eggs [24,25].

In this study, we aimed to determine the effects of coating quail table eggs with propolis, a natural product, on the interior egg quality characteristics during storage.

MATERIALS AND METHODS

This study was conducted at the Mustafa Kemal University Samandag Two-Year College in 2012. A total of 600 eggs were obtained from 12-wk-old Japonica quail (*Coturnix coturnix japonica*), which reached 90% egg production. Upon oviposition eggs were collected and brought to the laboratory for coating with various dosages of propolis extract.

The following treatments were included: control without any treatment (group I), coating with 5% (group II), 10% (group III), 15% (group IV) propolis extract, and coating with ethyl alcohol (group V).

The eggs in each treatment group were individually weighed in order to determine the egg weight loss along storage time. Before storage 20 fresh eggs (0 day storage) from each group were collected and analyzed within 2 h of being laid to determine the basic internal egg quality parameters. Following treatment, eggs were stored for 5 wk at room temperature (25°C). Every week, 20 eggs from each group were used to determine albumen pH, albumen height, albumen width, albumen length, yolk height, yolk length, and egg weight loss during storage.

Preparation of Propolis Solution

Propolis was collected from honey bees in Hatay Province, Turkey, in 2011 and extracted according to the method suggested by Krell [26; 25]. A 5% propolis solution was prepared by mixing 1,950 mL 70% ethanol and 50 g propolis, the 10% propolis solution was 1,900 mL 70% ethanol and 100 g propolis, and the 15% propolis solution was 1,850 mL 70% ethanol and 150 g of propolis. Solutions were kept in a sealed container and shaken twice daily for 2 wk.

Coating of Egg with Coating Solution and Alcohol

Eggs were immersed in the 5%, 10%, 15% propolis solutions and ethanol by hand for 1 min, and this process was repeated once more after 10 min. Following the second immersion the eggs were dried for 24 h at ambient temperature.

pH Measurement

After measurements for the albumen and yolk index parameters, the albumen was separated from the yolk. The volumes of thick and thin albumen were mixed with a spatula by hand before measuring the pH value. The pH of the albumen was measured using a pH 210 meter (Hanna Instruments, Woonsocket, RI).

Egg Weight Loss

Egg weight loss (EWL) during storage was calculated by individual measurement of initial and weekly weight changes in each group as follows:

$$\text{EWL (\%)} = (W_0 - W_1/W_0) \times 100$$

W_0 = Initial Egg Weight

W_1 = Egg Weight During Measurement Day

Albumen Index

Eggs from each treatment group were broken on a flat surface where the height of the albumen was measured in millimeters, halfway between the yolk and edge of the inner thick albumen, using a standard tripod micrometer. Measurements were taken from 3 points on the inner thick albumen at intervals of 10 mm from each other.

Measurement of albumen width was to 0.1 mm using a steel Vernier caliper. The albumen index (AI) was calculated as follows [27]:

$$\text{AI (\%)} = [\text{Albumen Height}/((\text{Albumen Length} + \text{Albumen Width})/2)] \times 100$$

Haugh Unit

Haugh unit (HU) values were calculated using the following formula [28]:

$$\text{HU} = 100 \text{Log}(H + 7.57 - 1.7 \times W^{0.37})$$

where H is the albumen height (millimeters) and W is the egg weight (grams).

Yolk Index

The yolk index (YI) was calculated as follows [29]:

$$\text{YI (\%)} = (\text{Yolk Height/Yolk Width}) \times 100$$

Data Analysis

In this study, the combined effect of propolis and storage time on internal egg qualities was evaluated. The data were subjected to a GLM using the *SPSS (SPSS for Windows release 13)* software package, with treatments and storage time as the main effects. When the main effects were significant at $P < 0.05$, differences between means were tested using Duncan's multiple range test.

RESULTS AND DISCUSSION

Albumen pH

To determine the changes in albumen pH during storage, it was necessary to measure the initial albumen pH immediately after oviposition time. Albumen quality was determined by albumen height and albumen pH. Initial albumen pH levels in each group increased with increasing storage time, but this increase in the control group and the groups coated with ethyl alcohol was higher than the increase in the other treatment groups coated with different concentrations of propolis (Table 1). The initial albumen pH level of the control group was 8.28, and that of the group coated with ethyl alcohol was 8.90. These values increased to 9.45 at the end of the fifth week (Table 1).

The difference in pH values at the beginning and end of the storage time in groups I, V, II, IV, and III were 0.55, 1.17, 0.30, 0.26, and 0.23, respectively (Table 1). The observed values supported previous studies by Caner [12] and Copur et al. [9], who found lower pH values at the end of storage when eggs were coated with chitosan, shellac, and propolis compared to uncoated eggs. In this study the albumen pH values recorded after the fifth week of storage were close to the pH value of quail eggs reported by Baylan [30], who found it to be 9.52 at 20°C after 30 d of storage,

Table 1. Effect of propolis coating on albumen pH during 5 wk of storage.

Group	Weeks					F	P
	0	1	2	3	4		
I	8.28 ± 0.12 ^d	9.00 ± 0.02 ^c	9.31 ± 0.03 ^{a,b}	9.32 ± 0.01 ^{a,b}	9.21 ± 0.01 ^b	9.45 ± 0.02 ^a	68.933
II	9.01 ± 0.02 ^c	8.98 ± 0.02 ^c	9.29 ± 0.01 ^{a,b}	9.29 ± 0.01 ^{a,b}	9.25 ± 0.01 ^b	9.31 ± 0.01 ^a	109.207
III	9.05 ± 0.01 ^d	8.97 ± 0.01 ^c	9.31 ± 0.01 ^b	9.35 ± 0.01 ^a	9.24 ± 0.01 ^c	9.28 ± 0.01 ^b	237.838
IV	9.01 ± 0.02 ^d	9.22 ± 0.01 ^c	9.31 ± 0.01 ^a	9.34 ± 0.01 ^a	9.19 ± 0.02 ^c	9.27 ± 0.02 ^b	68.381
V	8.90 ± 0.02 ^c	9.04 ± 0.01 ^d	9.24 ± 0.01 ^c	9.27 ± 0.01 ^b	9.22 ± 0.01 ^c	9.45 ± 0.01 ^a	172.272

I: Control group; II: 5% propolis; III: 10% propolis; IV: 15% propolis; V: alcohol.

^{a-c}Means with different superscripts within rows are significantly different ($P < 0.001$).

and by Aktan [31], who reported a pH of 9.78 after 7 d of storage.

Some researchers [3,5,7] have reported that the starting value of albumen pH for chicken egg was between 7.6 and 7.9. In the present experiment the basic albumen pH in quail egg was 8.84 (Table 6). This value is similar to that of previous studies, in which the initial albumen pH of quail egg was determined to be between 8.56 and 9.17 [30,32–33]. The pH of albumen increased with increasing storage time from 8.84 to 9.35 ($P < 0.001$) (Table 6). These results are in agreement with previous studies conducted by Silversides and Scott [3], Samli et al. [34], Caner [12], Akyürek and Okur [35], Aktan [31], Baylan et al. [30], Hristakieva et al. [36], and Jin et al. [37], who found an increase in the initial albumen pH level depending on storage time.

Egg Weight Loss

Internal egg quality characteristics are affected by factors such as egg surface coating with different materials and storage time. The egg weight loss of each treatment group increased significantly with storage time ($P < 0.001$) (Table 2) (Table 6). The egg weight loss of groups

I, II, III, IV, and V was 6.09, 5.72, 5.01, 4.67, and 5.81%, respectively. The highest egg weight loss values were obtained from the control group and samples coated with ethyl alcohol. While the egg weight loss in the first week was 1.16%, it reached 5.46% after the fifth week (Table 6). In the present experiment, egg weight loss differed significantly between all treatment groups ($P < 0.001$) (Table 7). The highest average egg weight loss was 3.57% in group I, followed by groups V (3.44%), II (3.27%), III (2.97%), and IV (2.17%).

These results are in agreement with those of Romao et al. [38], who reported egg weight losses of 2.72 and 2.39% during 2 wk of storage at 20°C in meat and laying type quails, respectively. Lacin et al. [39] also reported similar egg weight loss (2.56% during 2 wk of storage). In this study, the determined egg weight loss following 2 wk of storage was 2.18%, which was similar to the results given previously (Table 6). While in other studies the egg weight loss was 2.07% (32°C ambient temperature) [40] and 2.8% (at 32°C ambient temperature) [41] after 1 wk of storage, in the present experiment, the same losses were observed after 2 wk. Garip and Dere [42] also reported similar weight losses in quail eggs with 10 d of storage at 11°C, 21°C, and

Table 2. Effect of propolis coating on egg weight loss during 5 wk of storage (%).

Group	Weeks					F	P
	1	2	3	4	5		
I	1.28 ± 0.10 ^a	2.32 ± 0.14 ^a	3.64 ± 0.34 ^b	4.53 ± 0.56 ^b	6.09 ± 0.76 ^c	16.969	0.000
II	1.14 ± 0.06 ^d	2.30 ± 0.14 ^c	3.03 ± 0.21 ^c	4.29 ± 0.27 ^b	5.73 ± 0.46 ^a	46.002	0.000
III	1.08 ± 0.10 ^d	1.95 ± 0.13 ^{c,d}	2.75 ± 0.10 ^c	4.00 ± 0.59 ^b	5.01 ± 0.34 ^a	25.514	0.000
IV	1.05 ± 0.09 ^d	1.80 ± 0.11 ^c	2.47 ± 0.17 ^c	3.48 ± 0.33 ^b	4.68 ± 0.43 ^a	30.449	0.000
V	1.29 ± 0.12 ^d	2.54 ± 0.12 ^c	3.24 ± 0.32 ^c	4.35 ± 0.42 ^b	5.81 ± 0.32 ^a	35.700	0.000

I: Control group; II: 5% propolis; III: 10% propolis; IV: 15% propolis; V: alcohol.

^{a-d}Means with different superscripts within rows are significantly different ($P < 0.001$).

Table 3. Effect of propolis coating on value of albumen index during 5 wk of storage (%).

Group	Weeks						F	P
	0	1	2	3	4	5		
I	13.18 ± 0.54 ^a	10.40 ± 0.27 ^b	8.14 ± 0.43 ^c	7.20 ± 0.34 ^{c,d}	6.45 ± 0.29 ^{d,e}	5.65 ± 0.23 ^e	59.283	0.000
II	14.28 ± 0.38 ^a	10.72 ± 0.27 ^b	9.18 ± 0.29 ^c	7.53 ± 0.30 ^d	7.23 ± 0.30 ^d	6.34 ± 0.29 ^e	90.676	0.000
III	14.44 ± 0.39 ^a	10.40 ± 0.42 ^b	8.69 ± 0.33 ^c	7.87 ± 0.23 ^{c,d}	8.04 ± 0.26 ^{c,d}	7.29 ± 0.34 ^d	64.717	0.000
IV	14.29 ± 0.65 ^a	9.93 ± 0.32 ^b	9.03 ± 0.36 ^{b,c}	8.03 ± 0.29 ^{c,d}	7.89 ± 0.37 ^{c,d}	7.38 ± 0.21 ^d	42.128	0.000
V	14.05 ± 0.47 ^a	10.11 ± 0.42 ^b	8.04 ± 0.34 ^c	7.32 ± 0.28 ^{c,d}	6.67 ± 0.18 ^{d,e}	6.04 ± 0.15 ^e	82.932	0.000

I: Control group; II: 5% propolis; III: 10% propolis; IV: 15% propolis; V: alcohol.

^{a-e}Means with different superscripts within rows are significantly different ($P < 0.001$).

27°C. They found egg weight losses of 1.3, 3.1, and 3.7%, which were similar to the results of this study. In other experiments, egg weight losses of 1.58% were found in duck eggs stored for 11 d at 17°C [43], 1.66% in chicken eggs stored for 10 d at 21°C [34], and 2.60% in Fayumi chicken eggs stored for 8 d at 16°C and 75–80% relative humidity [44]. Differences in egg weight loss between studies may be due to different storage times, storage temperatures, egg sizes, or shell porosities.

In this study weight loss in uncoated eggs was found to be significantly higher than that found in coated eggs ($P < 0.001$) (Table 7). The samples with 15% propolis coating had significantly lower weight loss compared to the other two propolis concentrations. This finding is in agreement with the results reported by Aygun and Sert [25]. These authors reported 1.74, 1.75, and 1.37% and 2.86, 2.55, and 2.28% weight loss over 7 and 14 d of storage respectively for quail hatching eggs coated with 5, 10, and 15% propolis. Coating eggs using various concentrations of propolis may prolong their shelf life by preventing weight loss by limiting the loss of humidity and carbon dioxide through the shell.

Albumen Index

Some undesirable physicochemical changes in egg structure arose with increased storage times [34,45].

The albumen index significantly decreased with increasing storage time in all treatment groups ($P < 0.001$) (Table 3). The highest decrease in the albumen index after storage times compared with the initial albumen index values was in group V, followed by groups II, I, III, and IV. This result agreed with those of pre-

vious studies [4,9,12,35,37,46–48], which reported that the albumen index decreased with increasing storage time.

The index values with studies on quail were reported to be 12.56% by Salman and Tabeehk [49], 8.85% by Cabuk et al. [50], 14% by Shit et al. [51]; in 8-, 13-, 18-, and 23-wk-old quail (*Pharaoh coturnix*); those values were reported to be 10.0, 9.0, 9.0, and 13.0%, respectively, by Wilkanowska and Kokoszyński [33]; in Pharaoh and Manchurian Golden genotypes Genchev reported those values as 10.6% and 10.5%, respectively [52]. The difference in the initial albumen index value may be explained by the different genotypes and ages of quails used in previous studies.

The effect of storage time on the albumen index was significant ($P < 0.001$) (Table 6). In the present experiment the albumen index differed significantly among the treatment groups (Table 7). The highest albumen index values were obtained in groups II (9.24%), III (9.45%), and IV (9.45%), which were coated with propolis extract. The albumen index values of the control group and the group coated with ethyl alcohol were 8.50% and 8.71%, respectively.

Haugh Unit

The Haugh unit (HU) is related to albumen quality and measured as a function of the inner thick albumen height and egg weight. Weekly changes in HU values of all treatment groups during storage time are given in Table 4.

The basic HU value (95.28) decreased with increasing storage time. At the end of the storage time (week 5), the HU value was found to be 82.40. The differences between weeks were statistically significant ($P < 0.001$) (Table 6).

Table 4. Effect of propolis coating on Haugh units during 5 wk of storage.

Group	Weeks						F	P
	0	1	2	3	4	5		
I	92.92 ± 0.79 ^a	89.57 ± 0.46 ^b	83.90 ± 0.81 ^c	83.26 ± 0.72 ^c	80.78 ± 1.00 ^d	80.81 ± 0.56 ^d	44.540	0.000
II	95.50 ± 0.58 ^a	89.71 ± 0.52 ^b	87.76 ± 0.59 ^c	83.94 ± 0.63 ^d	83.67 ± 0.57 ^{d,e}	81.96 ± 0.80 ^e	65.429	0.000
III	96.19 ± 0.62 ^a	89.68 ± 0.87 ^b	86.70 ± 0.57 ^c	84.70 ± 0.70 ^d	85.17 ± 0.55 ^{e,d}	84.01 ± 0.67 ^d	48.531	0.000
IV	95.83 ± 1.32 ^{a,b}	89.30 ± 0.60 ^b	86.68 ± 0.68 ^c	85.33 ± 0.63 ^{c,d}	84.83 ± 0.63 ^{c,d}	84.24 ± 0.38 ^d	32.459	0.000
V	95.96 ± 0.58 ^a	89.76 ± 0.58 ^b	84.51 ± 0.82 ^c	82.63 ± 0.56 ^d	81.07 ± 0.46 ^d	81.00 ± 0.41 ^d	103.359	0.000

I: Control group; II: 5% propolis; III: 10% propolis; IV: 15% propolis; V: alcohol.

^{a-c}Means with different superscripts within rows are significantly different ($P < 0.001$).**Table 5.** Effect of propolis coating on yolk index during 5 wk of storage (%).

Group	Weeks						F	P
	0	1	2	3	4	5		
I	48.01 ± 1.04 ^a	40.64 ± 0.50 ^b	32.40 ± 0.59 ^c	32.31 ± 0.53 ^c	28.81 ± 0.41 ^d	26.93 ± 0.64 ^e	150.361	0.000
II	48.91 ± 0.66 ^a	40.79 ± 0.51 ^b	35.90 ± 0.57 ^c	32.74 ± 0.61 ^d	32.99 ± 0.76 ^d	27.00 ± 0.65 ^e	144.183	0.000
III	49.31 ± 0.62 ^a	41.07 ± 0.53 ^b	36.33 ± 0.60 ^c	33.43 ± 0.81 ^d	29.85 ± 0.54 ^e	27.95 ± 0.88 ^e	135.164	0.000
IV	49.78 ± 0.72 ^a	40.19 ± 0.48 ^b	36.14 ± 0.55 ^c	31.70 ± 0.56 ^d	28.92 ± 0.72 ^e	27.78 ± 0.69 ^e	175.610	0.000
V	49.58 ± 0.745 ^a	41.02 ± 0.50 ^b	34.85 ± 0.40 ^c	32.01 ± 0.73 ^d	29.37 ± 0.52 ^e	27.47 ± 0.47 ^f	206.191	0.000

I: Control group; II: 5% propolis; III: 10% propolis; IV: 15% propolis; V: alcohol.

^{a-f}Means with different superscripts within rows are significantly different ($P < 0.001$).**Table 6.** Mean weekly changes in initial egg quality for all treatments.

Properties	Weeks						F	P
	0	1	2	3	4	5		
Egg weight loss (g)		1.16 ± 0.04 ^e	2.18 ± 0.06 ^d	3.02 ± 0.11 ^c	4.13 ± 0.20 ^b	5.46 ± 0.22 ^a	128.842	0.000
Yolk index (%)	49.12 ± 0.34 ^a	40.74 ± 0.22 ^b	35.12 ± 0.28 ^c	32.44 ± 0.30 ^d	29.99 ± 0.31 ^e	27.43 ± 0.30 ^f	732.845	0.000
Albumen index (%)	14.05 ± 0.22 ^a	10.33 ± 0.15 ^b	8.61 ± 0.16 ^c	7.59 ± 0.13 ^d	7.25 ± 0.14 ^d	6.54 ± 0.13 ^e	301.661	0.000
Haugh unit	95.28 ± 0.38 ^a	89.60 ± 0.27 ^b	85.71 ± 0.33 ^c	83.93 ± 0.30 ^d	83.11 ± 0.35 ^{d,e}	82.40 ± 0.30 ^e	233.186	0.000
Albumen pH	8.84 ± 0.04 ^e	9.04 ± 0.01 ^d	9.29 ± 0.01 ^b	9.31 ± 0.05 ^{a,b}	9.22 ± 0.00 ^c	9.3 ± 0.01 ^a	132.549	0.000

I: Control group; II: 5% propolis; III: 10% propolis; IV: 15% propolis; V: alcohol.

^{a-f}Means with different superscripts within rows are significantly different ($P < 0.001$).**Table 7.** Means of interior egg quality at the end of the experiment on the different treatment groups.

Properties	Group					F	P
	I	II	III	IV	V		
Egg weight loss (g)	3.57 ± 0.26 ^a	3.27 ± 0.20 ^{a,b}	2.97 ± 0.20 ^{a,b}	2.67 ± 0.17 ^b	3.44 ± 0.20 ^a	3.084	0.001
Yolk index (%)	38.85 ± 0.72	36.47 ± 0.69	36.29 ± 0.72	35.82 ± 0.74	35.74 ± 0.74	0.769	0.546
Albumen index (%)	8.50 ± 0.28 ^b	9.24 ± 0.28 ^{a,b}	9.45 ± 0.26 ^a	9.45 ± 0.27 ^a	8.71 ± 0.29 ^{a,b}	2.564	0.037
Haugh unit	85.21 ± 0.51 ^c	86.97 ± 0.49 ^{a,b}	87.69 ± 0.47 ^a	87.73 ± 0.48 ^a	85.83 ± 0.56 ^{b,c}	5.071	0.001
Albumen pH	9.09 ± 0.03 ^b	9.19 ± 0.01 ^a	9.19 ± 0.01 ^a	9.22 ± 0.01 ^a	9.18 ± 0.02 ^a	4.762	0.001

I: Control group; II: 5% propolis; III: 10% propolis; IV: 15% propolis; V: alcohol.

^{a-c}Means with different superscripts within rows are significantly different ($P < 0.001$).

The HU values during 5 wk of storage were higher in groups IV (8.87) and III (87.69) than in groups II (86.97), V (85.83), and I (85.21) (Table 7). The decreasing HU values with increasing storage time were supported by previ-

ous studies [3,9,12,30,31,34–37] conducted with chicken and coturnix eggs. At the end of storage, the highest HU values were determined in groups IV and III. This result is in agreement with those of Allenoni and Antunes [53], Caner [12], and

Copur et al. [9], i.e., studies that found a higher HU value in coated eggs compared to uncoated eggs.

Yolk Index

The yolk index (YI) decreases during storage through the diffusion of water from the albumen to the yolk, resulting in changes in the vitellin membrane and in liquefaction of the yolk [3,54].

The YI decreased with increasing age of the hen eggs [55], though this trend was not confirmed in Japanese quail eggs, and the YI of quail eggs was higher than that of chicken eggs [56]. The differences in YI among treatment groups were insignificant ($P > 0.05$) (Table 7). The determined basic YI value (49.12%) was approximately similar to the values obtained by Zita et al. [57], who reported results of 49.11% for 9-wk-old quail, 47.66% for 13-wk-old quail, and 50.81% for 37-wk-old quail.

The YI values for fresh and good quality eggs were reported to be 43.67% by Kumar et al. [58] and Nowaczewski et al. [59], 45% by Dudusola [60], 46% by Canogullari et al. [61], 42.73% by Alkan et al. [62], and 47.80% by Zita et al. [57]. The effect of storage time on YI was statistically significant ($P < 0.001$) (Table 5).

The basic YI decreased significantly with increasing storage time (Table 6). The basic YI values decreased from 49.12 to 35.12, 32.44, 29.99, and 27.43% for 1-, 2-, 3-, 4-, and 5-wk storage times, respectively.

These results are in agreement with those of previous studies [4,9,12,30,44,59,63], which found a significant decrease in YI with increasing storage time in various poultry eggs.

CONCLUSIONS AND APPLICATIONS

1. Coating egg shell surfaces with propolis was found to be effective at protecting the internal egg quality parameters, and these parameters were negatively affected by storage time.
2. Coating eggs with propolis extract, a natural product, may help to ameliorate the decrease in quality during storage.

REFERENCES AND NOTES

1. Tayar, M. 2005. [http://mtayar.uludag.edu.tr/yumurtahijyeni.htm#Yumruta muhafaza yöntemleri](http://mtayar.uludag.edu.tr/yumurtahijyeni.htm#Yumruta%20muhafaza%20yontemleri), Erişim
2. Silversides, F. G., and P. Villeneuve. 1994. Is the Haugh unit correction for egg weight valid for eggs stored at room temperature? *Poult. Sci.* 73:50–55.
3. Silversides, F. G., and T. A. Scott. 2001. Effect of storage and layer age on quality of eggs from two different lines of hens. *Poult. Sci.* 80:1240–1245.
4. Tilki, M., and M. Saatci. 2004. Effects of storage time on external and internal characteristics in partridge (*Alectoris graeca*) eggs. *Revue Méd. Vét.* 155:561–564.
5. Li-Chan, E. C. Y., W. D. Powrie, and S. Nakai. 1994. The Chemistry of Eggs and Egg Products. Page 118 in *Egg Science and Technology*. 4th ed. Stadelman, W. J., and O. J. Cotterill, eds. The Hawort Press, Binghamton, NY. 13904–1580.
6. Heath, J. L. 1977. Chemical and related osmotic changes in egg albumen during storage. *Poult. Sci.* 56:822–828.
7. French, N. A., and S. G. Tullet. 1991. Variation in the eggs of poultry species. Pages 59–77. *Avian Incubation 22nd Poultry Science Symposium*, Butterworth-Heinemann Ltd., London.
8. Wong, Y. G., T. J. Herald, and K. A. Hachmeister. 1996. Evaluation of mechanical and barrier properties of protein coating on eggshell. *Poult. Sci.* 75:417–422.
9. Copur, G., O. Camci, N. Sahinler, and A. Gul. 2008. The effect of propolis egg shell coatings on interior egg quality. *Arch. Geflügelk.* 72:35–40.
10. Bhale, S., H. K. No, N. Prinyawiwatkul, A. J. Faar, K. Nadarajah, and S. P. Meyers. 2003. Chitosan coating improves shell life of eggs. *J. Food Sci.* 68:2378–2383.
11. Hisil, Y., and S. Otles. 1997. Changes of vitamin B-1 concentrations during storage of hen eggs. *Food Sci. Technol.-Lab.* 30:320–323.
12. Caner, C. 2005. The effect of edible eggshell coatings on egg quality and consumer perception. *J. Sci. Food Agric.* 85:1897–1902.
13. Greenaway, W., T. Scaysbrook, and F. R. Whatley. 1990. The composition and plant origins of propolis: A report of work at Oxford. *Bee World.* 71:107–118.
14. Schmidt, J. O. 1997. Bee products: Chemical composition and application. Pages 15–26 in *Bee Products: Properties, Applications, and Apitherapy*. Mizrahi, A., and Y. Lensky, eds. Plenum Press, New York and London.
15. Yamamoto, T. 1997. Present state of basic studies on propolis in Japan. Pages 107–120 in *Bee Products: Properties, Applications, and Apitherapy*. Mizrahi, A., and Y. Lensky eds., Plenum Press, New York and London.
16. Burdock, G. A. 1998. Review of the biological properties and toxicity of bee propolis (propolis). *Food. Chem. Toxicol.* 36:347–363.
17. Hegazi, A. G., F. K. Abd El Hady, and F. A. Abd Allah. 2000. Chemical composition and antimicrobial activity of European propolis. *Z. Naturforsch.* 55:70–75.
18. Nagai, T., M. Sakai, H. Inoue, and N. Suzuki. 2001. Antioxidative activities of some commercially honeys, royal jelly and propolis. *Food Chem.* 75:237–240.

19. Castaldo, S., and F. Capasso. 2002. Propolis, an old remedy used in modern medicine. *Fitoterapia*. 73:S1–S6.
20. Kartal, M., S. Yildiz, S. Kaya, S. Kurucu, and G. Topcu. 2003. Antimicrobial activity of propolis samples from two different regions of Anatolia. *J. Ethnopharmacol.* 86:69–73.
21. Orsolic, N., and I. Basic. 2003. Immunomodulation by water soluble derivative of propolis a factor of antitumor reactivity. *J. Ethnopharmacol.* 84:265–273.
22. Torre, A., G. Imbroglini, and M. Guccione. 1990. Preliminary observations on the action of propolis based preparations against *Botrytis cinerea* Pers. on strawberries. *Apicoltura*. 6:169–177.
23. Ozdemir, A. E., E. Erturk, N. Sahinler, M. Kaplankiran, and A. Gul. 2005. Propolis uygulamalarının fremont mandalinlerin muhafazasına etkileri. III. Bahce Urunlerinde Muhafaza ve Pazarlama Sempozyumu. 09–09 Eylül 2005. Hatay. 204–211.
24. Aygun, A., D. Sert, and G. Copur. 2012. Effects of propolis on eggshell microbial activity, hatchability, chick performance in Japanese quail (*Coturnix coturnix japonica*) eggs. *Poult. Sci.* 91:1018–1025.
25. Aygun, A., and Durmus Sert. 2013. Effects of prestorage application of propolis and storage time on eggshell microbial activity, hatchability, and chick performance in Japanese quail (*Coturnix coturnix japonica*) eggs. *Poult. Sci.* 92:3330–3337.
26. Krell, R. 1996. Value-added products from beekeeping. FAO Agricultural Services Bulletin No. 124. Food and Agriculture Organization of the United Nations, Rome.
27. Heiman, V., and J. S. Carver. 1936. The albumen index as a physical measurement of observed egg quality. *Poult. Sci.* 15:141–148.
28. Haugh, R. R. 1937. A new method for determining the quality of an egg. *US Egg Poult.* 39:27–49.
29. Funk, E. M. 1948. The relation of the yolk index determined in natural position to the yolk index, as determined after separating the yolk from the albumen. *Poult. Sci.* 27:367.
30. Baylan, M., S. Canogullari, T. Ayasan, and G. Copur. 2011. Effects of dietary selenium source, storage time, and temperature on the quality of quail eggs. *Biol. Trace Elem. Res.* 143:957–964.
31. Aktan, S. 2011. Effects of age and storage duration on relationships among albumen quality traits and egg weight in Japanese quails. *J. Anim. Vet. Adv.* 10:3340–3344.
32. Teuşan, A., V. Teuşan, and A. A. Prelipcean. 2009. Research regarding some physical quality values of Japanese quail eggs obtained at the middle of the laying stage. *Lucrări Stiinţifice Zootehnie si Biotehnologii.* 42:446–451.
33. Wilkanowska, A., and D. Kokoszynski. 2012. Layer age and quality of pharaoh quail eggs. *J. Central Eur. Agric.* 13:10–21.
34. Samli, H. E., A. Agma, and N. Senkoylu. 2005. Effects of storage time and temperature on egg quality in old laying hens. *J. Appl. Poult. Res.* 14:548–553.
35. Akyurek, H., and A. Agma Okur. 2009. Effect of storage time, temperature and hen age on egg quality in free-range layer hens. *J. Anim. Vet. Adv.* 8:1953–1958.
36. Hristakieva, P., M. Lalev, M. Oblakova, N. Mincheva, and Ivanova. 2011. Effect of storage duration on the quality of hatching turkey eggs. *Arch. Zootechnica* 14:57–65.
37. Jin, Y. H., K. T. Lee, W. I. Lee, and Y. K. Han. 2011. Effects of storage temperature and time on the quality of eggs from laying hens at peak production. *Asian-Aust. J. Anim. Sci.* 24:279–284.
38. Romao, J. M., T. G. V. Moraes, R. S. C. Teixeira, W. M. Cardoso, and C. C. Buxade. 2008. Effect of egg storage length on hatchability and weight loss in incubation of egg and meat type Japanese quails. *Braz. J. Poult. Sci.* 10:143–147.
39. Lacin, E., O. Coban, and N. Sabuncuoglu. 2008. Effects of egg storage material and storage period on hatchability in Japanese quail. *Asian-Aust. J. Anim. Sci.* 21:1183–1188.
40. Romao, J. M., T. G. V. Moraes, E. E. Silva, R. S. C. Teixeira, and W. M. Cardoso. 2010. Incubation of Japanese quail eggs stored at tropical temperatures. *Livestock Research for Rural Development*. 22(1). Article 2. Available from: <http://www.lrrd.org/lrrd22/1/roma22002.htm>. Accessed on January 20, 2015.
41. Dudusola, I. O. 2009. Effects of storage methods and length of storage on some quality parameters of Japanese quail eggs. *Tropicultura*. 27:45–48.
42. Garip, M., and S. Dere. 2011. The effect of storage period and temperature on weight loss in quail eggs and the hatching weight of quail chicks. *J. Anim. Vet. Adv.* 10:2363–2367.
43. Onbasilar, E. E., O. Poyraz, and E. Erdem. 2007. Effects of egg storage period on hatching egg quality, hatchability, chick quality and relative growth in Pekin ducks. *Arch. Geflügelk.* 71:187–191.
44. Khan, M. J. A., S. H. Khan, A. Bukhsh, M. I. Abbass, and M. Javed. 2013. Effect of different storage period on egg weight, internal egg quality and hatchability characteristics of Fayumi eggs. *Ital. J. Anim. Sci.* 12:323–328.
45. Jones, D. R., and M. T. Musgrove. 2005. Effects of extended storage on egg quality factors. *Poult. Sci.* 84:1774–1777.
46. Yilmaz, A. A., and Z. Bozkurt. 2009. Effects of hen age, storage period and stretch film packaging on internal and external quality traits of table eggs. *Lucrări Stiinţifice Zootehnie si Biotehnologii.* 42:462–469.
47. Demirel, S., and K. Kirikci. 2009. Effect of different egg storage times on some egg quality characteristics and hatchability of pheasants (*Phasianus colchicus*). *Poult. Sci.* 88:440–444.
48. Nowaczewski, S., H. Kontecka, A. Rosiński, S. Koberling, and P. Koronowski. 2010. Egg quality of Japanese quail depends on layer age and storage time. *Folia Biol. (Kraków)*. 58:201–207.
49. Salman, M. A., and A. Tabeekh. 2011. Evaluation of some external and internal egg quality traits of quails reared in Basrah city. *Bas. J. Vet. Res.* 10:78–84.
50. Cabuk, M., S. Eratak, A. Alcicek, and M. Bozkurt. 2014. Effects of herbal essential oil mixture as a dietary supplement on egg production in quail. *Sci. World J.* 2014:573470, published online 2014 January 23. Doi:10.1155/2014/573470.
51. Shit, N., R. P. Singh, K. V. H. Sastry, R. Agarwal, R. Singh, N. K. Pandey, and J. Mohan. 2012. Effect of dietary L-ascorbic acid (L-AA) on production performance, egg quality traits and fertility in Japanese quail (*Coturnix coturnix japonica*) at low ambient temperature. *Asian-Aust. J. Anim. Sci.* 25:1009–1014.
52. Genchev, A. 2012. Quality and composition of Japanese quail eggs (*Coturnix Japonica*). *Trakia J. Sci.* 10:91–101.

53. Allenoni, A. C. C., and A. J. Antunes. 2004. Internal quality of eggs coated with whey protein concentrate. *Sci. Agric. (Piracicaba, Braz.)* 61:276–280.
54. Scott, T. A., and F. G. Silversides. 2000. The effect of storage and strain of hen on egg quality. *Poult. Sci.* 79:1725–1729.
55. Zita, L., E. Tumova, and L. Stolc. 2009. Effects of genotype, age and their interaction on egg quality in brown-egg laying hens. *Acta Vet. Brno.* 78:85–91.
56. Zita, L., L. Zdenek, E. Tumova, and L. Klesalova. 2012. Technological quality of eggs in relation to the age of laying hens and Japanese quails. *R. Bras. Zootec.* 41:2079–2084.
57. Zita, L., L. Zdeněk, and K. Ludmila. 2013. The effect of the age of Japanese quails on certain egg quality traits and relationships. *Vet. Arhiv.* 83:223–232.
58. Kumari, P. B., G. B. Ramesh, G. M. Prakash, and R. A. Rajasekhar. 2008. A Study on egg quality traits in japanese quails (*Coturnix Coturnix Japonica*). *Tamilnadu J. Vet. Anim. Sci.* 4:227–231.
59. Nowaczewski, S., H. Kontecka, G. Elminowska-Wenda, A. Rosiński, M. Bednarczyk, and A. Kucharska. 2010. Egg quality traits in Japanese quail divergently selected for yolk cholesterol level. *Arch. Geflügelk.* 74:141–144.
60. Dudusola, I. O. 2010. Comparative evaluation of internal and external qualities of eggs from quail and guinea fowl. *Int. Res. J. Plant Sci.* 1:112–115.
61. Canogullari, S., M. Baylan, Z. Erdogan, V. Duzguner, and A. Kucukgul. 2010. The effects of dietary garlic powder on performance, egg yolk and serum cholesterol concentrations in laying quails. *Czech J. Anim. Sci.* 55: 286–293.
62. Alkan, S., K. Karabag, A. Galic, T. Karsli, and M. S. Balcioglu. 2010. Effects of selection for body weight and egg production on egg quality traits in japanese quails (*Coturnix coturnix japonica*) of different lines and relationships between these traits. *Kafkas Univ. Vet. Fak. Derg.* 16:239–244.
63. Khan, M. J. A., S. H. Khan, A. Bukhsh, and M. Amin. 2014. The effect of storage time on egg quality and hatchability characteristics of Rhode Island Red (RIR) hens. *Vet. Arhiv.* 84:291–303.

Acknowledgments

This study was supported by the Scientific Research Project Unit of Mustafa Kemal University, Turkey (Project number: 1201M0110).