

Table 1. Patient characteristics and adverse events by treatment group

	Anti-TNF	JAKi
Mean age start of treatment (SD), yrs	50.8 (12.6)	57.6 (11.9)
Female, n (%)	3122 (58.8)	392 (79.5)
Disease duration, median (IQR)	7.0 (2.7-13.7)	9.9 (4.9-16.8)
First line biologic, n (%)	2614 (49.3)	117 (23.7)
Rheumatoid Arthritis	1385 (41.1)	339 (95.2)
Ankylosing Spondylitis	1031 (30.6)	1 (0.3)
Psoriatic Arthritis	957 (28.4)	16 (4.5)
DAS28-ESR	4.3 (1.4)	4.7 (1.4)
Survival first year (IC 95%)	73.3 (71.9-74.6)	69.7 (66.0-73.0)
Charlson Index, mean (SD)	1.9 (1.3)	2.3 (1.6)
Reason to stop therapy (n)*:		
Lack of efficacy	1534 (53.2)	57 (58.8)
Adverse event	723 (25.1)	33 (34.0)
Adverse events (AE)*		
Serious infections	14.2 (12.4-16.2)	33.2 (19.3-57.3)
Herpes zoster	5.7 (4.6-7.1)	12.8 (5.3-30.7)
Tuberculosis	0.7 (0.4-1.3)	0.0 (0.0-0.0)
Malignancy/Neoplasia	10.2 (8.7-11.9)	15.3 (6.9-34.2)
Cardiac events	13.9 (12.2-16.0)	30.7 (17.4-54.0)
GI perforation	1.2 (0.8-1.9)	10.2 (3.8-27.3)
Vascular events	9.8 (8.3-11.5)	25.6 (13.8-47.5)

*Data show the incidence rate ratio per 1000 patient-years (PYs; 95% CI)

Conclusion: Serious infections and herpes zoster tend to be more frequent in patients on JAKi. However patients on JAKi were older, presented higher comorbidity and have a longer disease duration.

Disclosure of Interests: None declared

DOI: 10.1136/annrheumdis-2021-eular.2645

OP0124

EFFECTS OF NINTEDANIB IN PATIENTS WITH PROGRESSIVE FIBROSING INTERSTITIAL LUNG DISEASE ASSOCIATED WITH RHEUMATOID ARTHRITIS (RA-ILD) IN THE INBUILD TRIAL

C. Kelly¹, E. Matteson², M. Aringer³, G. R. Burmester⁴, H. Mueller⁵, L. Moros⁶, K. Rohr⁶, M. Kolb⁷ on behalf of the INBUILD Trial Investigators.. ¹Newcastle University, Institute of Cellular Medicine, Newcastle upon Tyne, United Kingdom; ²Mayo Clinic College of Medicine and Science, Division of Rheumatology, Rochester, United States of America; ³University Medical Center & Faculty of Medicine, TU Dresden, Rheumatology, Medicine III, Dresden, Germany; ⁴Charité-University Medicine Berlin, Department of Rheumatology and Clinical Immunology, Berlin, Germany; ⁵Boehringer Ingelheim Pharma GmbH & Co. KG, Biostatistics and Data Sciences, Biberach, Germany; ⁶Boehringer Ingelheim International GmbH, TA Inflammation Med, Ingelheim am Rhein, Germany; ⁷McMaster University and St. Joseph's Healthcare, Department of Medicine, Hamilton, Canada

Background: In the INBUILD trial in subjects with progressive fibrosing ILDs other than idiopathic pulmonary fibrosis (IPF), nintedanib reduced the rate of decline in FVC (mL/year) over 52 weeks by 57% compared with placebo.

Objectives: To assess the rate of decline in FVC in subjects with RA-ILD in the INBUILD trial.


Methods: Subjects in the INBUILD trial had a chronic fibrosing ILD other than IPF, reticular abnormality with traction bronchiectasis (with or without honeycombing) of >10% extent on high-resolution computed tomography (HRCT), forced vital capacity (FVC) ≥45% predicted, diffusing capacity of the lungs for carbon monoxide ≥30%–<80% predicted, and met criteria for progression of ILD within the 24 months before screening, despite management deemed appropriate in clinical practice. Patients taking stable doses of approved medications to treat RA or connective tissue disease could participate, except that the protocol excluded those taking azathioprine, cyclosporine, mycophenolate mofetil, tacrolimus, rituximab, cyclophosphamide, or oral glucocorticoids >20 mg/day. We analysed the rate of decline in FVC (mL/year) over 52 weeks and adverse events in subjects with RA-ILD.

Results: Of 663 subjects who received trial medication, 89 had RA-ILD (42 nintedanib, 47 placebo), of whom 60.7% were male, 64.0% were current or former smokers, 86.5% had a usual interstitial pneumonia (UIP)-like pattern on HRCT; 93.3% had received confirmation of their RA diagnosis from a rheumatologist. At baseline, 21.3% of subjects were taking biologic disease-modifying anti-rheumatic drugs (DMARDs), 53.9% were taking non-biologic DMARDs and 73.0% were taking glucocorticoids (≤20 mg/day prednisone or

equivalent). At baseline, mean (SD) age was 66.9 (9.6) years, time since RA diagnosis was 9.9 (9.4) years, time since ILD diagnosis was 3.6 (3.2) years, FVC was 71.5 (16.2) % predicted and C-reactive protein was 13.7 (22.5) mg/L. The adjusted mean (SE) rate of decline in FVC over 52 weeks was -82.6 (41.3) mL/year in the nintedanib group versus -199.3 (36.2) mL/year in the placebo group (difference 116.7 mL/year [95% CI 7.4, 226.1]; nominal p=0.037), consistent with findings in the overall trial population (Figure). As in the overall trial population, the most common adverse event in subjects with RA-ILD was diarrhoea (reported in 54.8% of the nintedanib group and 25.5% of the placebo group). Adverse events led to permanent discontinuation of trial drug in 19.0% of subjects in the nintedanib group and 12.8% of subjects in the placebo group.

Conclusion: In the INBUILD trial, nintedanib slowed the rate of decline in FVC in patients with progressive fibrosing RA-ILD, with adverse events that were manageable for most patients. The efficacy and safety of nintedanib in subjects with RA-ILD were consistent with those observed in the overall trial population.

Figure. Rate of decline in FVC (mL/year) over 52 weeks with nintedanib versus placebo in all subjects and in subjects with RA-ILD in the INBUILD trial.



Acknowledgements: The INBUILD trial was funded by Boehringer Ingelheim. Medical writing support was provided by Fleishman Hillard Fishburn, London, UK. The authors meet criteria for authorship as recommended by the International Committee of Medical Journal Editors (ICMJE).

Disclosure of Interests: Clive Kelly Speakers bureau: Boehringer Ingelheim, Consultant of: Boehringer Ingelheim, Eric Matteson Speakers bureau: Boehringer Ingelheim, Consultant of: Boehringer Ingelheim and Gilead Sciences, Grant/research support from: Sun Pharmaceuticals and Pfizer, Martin Aringer Speakers bureau: AbbVie, AstraZeneca, Bristol-Myers Squibb, Boehringer Ingelheim, Chugai, Gilead, GlaxoSmithKline, HEXAL, Lilly, MSD, Novartis, Pfizer, Roche, Sanofi and UCB, Consultant of: AbbVie, AstraZeneca, Bristol-Myers Squibb, Boehringer Ingelheim, GlaxoSmithKline, Lilly, Merck Sharp & Dohme, Novartis, Pfizer, Roche, Sanofi and UCB, Gerd Rüdiger Burmester Speakers bureau: AbbVie, Bristol-Myers Squibb, Boehringer Ingelheim, Gilead, Lilly, Merck Sharp & Dohme, Pfizer, Roche and Sanofi, Heiko Mueller Employee of: Currently an employee of Boehringer Ingelheim, Lizette Moros Employee of: Currently an employee of Boehringer Ingelheim, Klaus Rohr Employee of: Currently an employee of Boehringer Ingelheim, Martin Kolb Consultant of: Algernon, Boehringer Ingelheim, Pieris Pharmaceuticals and Roche, Grant/research support from: Boehringer Ingelheim, Pieris Pharmaceuticals and Prometic

DOI: 10.1136/annrheumdis-2021-eular.969

OP0125

TOFACITINIB IN RHEUMATOID ARTHRITIS-ASSOCIATED INTERSTITIAL LUNG DISEASE: EFFICACY AND SAFETY ANALYSIS FROM TREASURE REAL-LIFE DATA

U. Kalyoncu¹, E. Bilgin¹, A. Erden², H. Satış³, A. Tufan³, E. Tekgoz⁴, A. Ateş⁵, B. N. Coşkun⁶, B. Yağız⁶, O. Küçükşahin⁷, V. Yazısız⁸, G. Kimyon⁹, C. Bes¹⁰, A. İ. Ertenli¹, S. Kiraz¹. ¹Hacettepe University, Internal Medicine, Rheumatology, Ankara, Turkey; ²Ankara City Hospital, Internal Medicine, Rheumatology, Ankara, Turkey; ³Gazi University, Internal Medicine, Rheumatology, Ankara, Turkey; ⁴University of Health Sciences Turkey, Gulhane Faculty of Medicine, Internal Medicine, Rheumatology, Ankara, Turkey; ⁵Ankara University, Internal Medicine, Rheumatology, Ankara, Turkey; ⁶Uludağ University, Internal Medicine, Rheumatology, Bursa, Turkey; ⁷Yıldırım Beyazıt University, Internal Medicine, Rheumatology, Ankara, Turkey; ⁸Akdeniz University, Internal Medicine, Rheumatology, Antalya, Turkey; ⁹Mustafa Kemal University,

Objectives: To assess long-term safety across the FIL program regarding infections, including serious infections (SI).