

The Relationship between an Accessory Maxillary Ostium and Variations in Structures Adjacent to the Maxillary Sinus without Polyps

Gozde Serindere¹ Kaan Gunduz² Hakan Avsever³

¹Department of Dentomaxillofacial Radiology, Faculty of Dentistry, Hatay Mustafa Kemal University, Hatay, Turkey.

²Department of Dentomaxillofacial Radiology, Faculty of Dentistry, Ondokuz Mayıs University, Samsun, Turkey.

³Department of Dentomaxillofacial Radiology, Gulhane Faculty of Dentistry, Health Sciences University, Ankara, Turkey.

Address for correspondence Gozde Serindere, associate professor, Department of Dentomaxillofacial Radiology, Faculty of Dentistry, Hatay Mustafa Kemal University, 31060 Hatay, Turkey (e-mail: gozdeserindere@mku.edu.tr).

Int Arch Otorhinolaryngol 2022;26(4):e548–e555.

Abstract

Introduction The maxillary sinus and its variations are very important to dentistry and rhinology.

Objective To investigate the effect of the accessory maxillary ostium (AMO) on the variations of adjacent structures of the maxillary sinus.

Methods The computed tomography (CT) images of 400 patients were retrospectively evaluated. The prevalence of AMO was calculated. The relationship between morphological variations of adjacent structures of maxillary sinus such as agger nasi cell (ANC), Haller cell (HC), nasal septum deviation (NSD), hypertrophy of inferior concha (HIC), pneumatization of middle concha (PMC), mucus retention cyst (MRC), mucosal thickening (MT), and maxillary sinusitis (MS), as well as the presence of AMO, were investigated.

Results Presence of AMO was diagnosed in 42 patients (10.5%), having been found in 4.5% of the patients only on the right side, in 1.25% of the patients only on the left side, and in 4.75% of the patients on both sides. There is an increasing incidence of ANC, HC, NSD, HIC, and PMC in the presence of AMO and MS. There is a decreasing incidence of MRC in the presence of AMO. Furthermore, AMO does not affect the incidence of MT.

Conclusion This study showed that most parameters, except for MRC and MT, had increasing incidence in the presence of AMO. It is important for radiologists and rhinologists to have knowledge about the location of AMO and the presence of variations of MS adjacent structures to avoid surgical complications.

Keywords

- ▶ computed tomography
- ▶ paranasal sinuses
- ▶ sinus anatomy
- ▶ sinusitis

Introduction

The natural ostium of the maxillary sinus is anteriorly placed and has a transversely oval form. It is not visible with nasal

endoscopic evaluation.¹ The ostium of the maxillary sinus is at the highest part of the medial wall of the sinus and is, therefore, poorly placed in terms of free drainage; in addition, it does not open into the nasal fossa, but into the narrow

received
May 18, 2021
accepted after revision
October 17, 2021
published online
January 28, 2022

DOI <https://doi.org/10.1055/s-0042-1742325>.
ISSN 1809-9777.

© 2022. Fundação Otorrinolaringologia. All rights reserved.
This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

ethmoidal infundibulum, and its inflammation may further inhibit drainage.²

In addition to a natural ostium connecting the maxillary sinus to the middle meatus, endoscopic *in vivo* evaluations commonly reveal the presence of an accessory maxillary ostium (AMO) in the maxillary fontanelle.³ The AMO is generally situated on the posterior fontanelle of the lateral nasal wall. The maxillary hiatus should not be confused with AMO.¹ An AMO is observed in 30% of patients suffering from chronic maxillary sinusitis (MS) and in 10 to 20% of healthy individuals.^{4,5}

Chronic sinusitis causes significant morbidity despite appropriate medical and surgical treatment. Anatomical variations, including AMO, play a role in the pathogenesis of chronic MS. An AMO impairs mucociliary clearance of the maxillary sinus owing to the recycling of mucus between the natural and accessory ostia, and this process may result in chronic MS.⁶ It is not clear whether an AMO is congenital or acquired. Some authors claim that it may develop after acute MS.⁷

The aim of this study was to evaluate the incidence of AMO and the relationship between an AMO and variations in structures adjacent to the maxillary sinus.

Materials and Methods

The present study is a retrospective study, approved by the Local Ethics Committee of the Faculty of Medicine of the Hatay Mustafa Kemal University (decision date: 26/12/2019, decision number: 07). The medical history of all patients was noted through the picture archiving and communication system (PACS) of the university hospital. The computed tomography (CT) scans of 400 patients (male = 222, female = 178) who had been referred to the Department of Dento-maxillofacial Radiology of our University Hospital were evaluated.

The exclusion criteria from this study were:

- Patients with maxillofacial trauma.
- Patients undergoing sinus surgery.
- Patients diagnosed with nasal polyps, acute sinusitis, inverted papilloma, choanal atresia, or advanced nasal septal deviation (NSD) contacting the lateral nasal wall.
- CT scans with low-quality images.

The study of Yenigun et al.⁸ investigated the prevalence of AMO and the relationship between an AMO and anatomical variations in the adjacent anatomical structures, namely the agger nasi cell (ANC), Haller cells (HC), NSD, pneumatization of the middle turbinate or concha (concha bullosa; PMC), hypertrophy of the inferior concha (HIC), maxillary sinusitis (MS), mucus retention cysts (MRC), and mucosal thickening (MT) (–Figs. 1,2,3,4,5). The following definitions were taken into account in the evaluation of the parameters:

ANC: Agger nasi cells are frontal ethmoidal cells, located in front of the anterior end of the middle concha and uncinate process in the nasal lateral wall.⁹

HC: Haller cells are alternatively referred to as infraorbital ethmoid cells, as they arise from anterior ethmoid cells and are located in the medial orbital floor.¹⁰




Fig. 1 Schematic diagram of the accessory maxillary ostium.

NSD: Nasal septal deviation is a common condition in which the bone or cartilage of the nasal septum is deviated from the midline of the face.¹¹

PMC: Pneumatization of the middle turbinate, or concha, is caused by an air-filled cavity in a concha or turbinate.¹²

HIC: Hypertrophy of the inferior concha is a condition encountered with NSD, caused by the increase in the size of the inferior concha (also called compensatory hypertrophy of the inferior concha). The hypertrophy is observed on the opposite side to the NSD. For example, if the nasal septum deviates to the right, the hypertrophy is observed to the left.¹³

MS: Inflammation of the maxillary sinuses. The symptoms of sinusitis are headache, generally in the region of the non-healthy sinus, and possibly a foul-smelling nasal or pharyngeal discharge, sometimes combined with systemic infection such as fever and malaise. The skin over the non-healthy sinus may be tender, hot or even red-dened because of inflammation. In radiological examination, opacification (or clouding) of the translucent sinus is generally observed because of mucus involvement.¹⁴

MRC: According to the study of Bhattacharyya,¹⁵ a diagnosis of MRC is based on the following criteria:

- a homogeneous, dome-shaped cyst with sharp borders;
- a lack of bony destruction;
- a lack of communication with tooth roots;
- a smooth, spherical contour along the cyst-free border.

MT: Mucosal thickening often coexists with chronic MS,¹⁶ and is probably pathological when larger than 2 mm. However, even MT up to 4 to 5 mm can be asymptomatic and go unnoticed by the patient.¹⁷

A Toshiba Aquilion (Canon Medical Systems Corp. Ōtawara, Tochigi, Japan) CT scanner was used for all the maxillofacial CT procedures. A routine maxillofacial CT protocol was conducted. All evaluations were performed with a Lenovo IdeaPad 520 notebook (Lenovo Ltda. Hong Kong,




Fig. 2 Agger nasi cell (A and B – white arrows) and accessory maxillary ostium (C and D – blue arrows).

China) with 15.6-inch and 1920 × 1080 resolution. All evaluations were done by a single blinded observer, Gozde Serindere, who had 6 years of clinical experience. In case there was any disagreement, these were solved by consensus with Professor Kaan Gunduz, who had nearly 15 years of clinical experience.

During the evaluations, the data of each patient were recorded in an Excel table (Microsoft Co. Redmond, Washington, USA). The Statistical Package Social Sciences (SPSS, StataCorp LLC. College Station, Texas, USA) software, version 16, was used to describe and analyze the data. The data were also analyzed by odds ratio (OR), in such way that the OR was obtained for the presence or absence of an AMO, as compared with the presence or absence of each of the potential complications. The significance level was set at $p = 0.05$.

Results

Of the 400 patients included in the study, 222 were male (55.5%) and 178 were female (44.5%). The patients' ages were between 7 and 87 years, and their mean age was of 40.7 ± 18.3 years.

The Incidence of AMO

In this study, the presence of an AMO and accompanying morphological variations in neighboring structures were assessed in 800 sides of 400 patients. We diagnosed 42




Fig. 3 A – Haller cell (white arrows); B – Big Haller cell (blue arrow).

patients (10.5%) with AMO, having been found in 4.5% of the patients on the right side, in 1.25% of the patients on the left side, and bilaterally in 4.75% of the patients.




Fig. 4 A – Bilateral pneumatization of middle concha and left nasal septum deviation B – Right nasal septum deviation with left hypertrophy of inferior concha.

The Relationship between AMO and ANC

We identified ANC in 49.3% of the studied subjects. In 31% (26/84) of the cases, ANC was diagnosed along with an AMO, 28.6% of which (12/42) the AMO was on the right side, 2.4% (1/42) it was on the left side, and 15.5% (13/84) where it was bilateral. Among the 716 sides in 358 patients without an AMO, ANC was found in 2.2% (16/716) of the cases, with 1.7% (6/358) on the right side, 1.1% (4/358) on the left side, and 0.8% (6/716) bilateral. The incidence of ANC increased by 77% in the presence of an AMO (OR = 1.77) (►Table 1).

The Relationship between AMO and HC

We identified HC in 10.3% of the studied subjects. In 5.9% (5/84) of the cases, it was found along with an AMO, 2.4% (1/42) of which the AMO was on the right side, and 4.7% (4/84) where it was bilateral. In patients without an AMO, HC was found in 5.16% (37/716), with 4.75% (17/358) on the right side, 1.4% (5/358) on the left side, and bilaterally in 2.1% (15/716). The incidence of HC increased by 20% in the presence of an AMO (OR = 1.20) (►Table 1).

The Relationship between AMO and NSD

We identified NSD in 41.5% of the studied subjects. In 23.8% (20/84) of the cases, it was found along with an AMO, 23.8% (10/42) of which it was on the right side, and 11.9% (10/84) it was bilateral. In patients without an AMO, NSD was found in 3.07% (22/716) of the cases, with 2.2% (8/358) on the right side, 1.4% (5/358) on the left side, and 1.25% (9/716) bilateral. The incidence of NSD increased by 32% in the presence of an AMO (OR = 1.32) (►Table 1).

The Relationship between AMO and HIC

We identified HIC in 35% of the studied subjects. It was found in 22.6% (19/84) of the cases with an AMO, with 23.8% (10/42) being on the right side and 10.7% (9/84) bilateral. In patients without an AMO, HIC was found in 3.2% (23/716) of the cases, with 2.2% (8/358) on the right side, 1.4% (5/358) on the left side, and 1.4% (10/716) bilateral. The incidence of HIC increased by 61% in the presence of an AMO (OR = 1.61) (►Table 1).

The Relationship between AMO and PMC

We identified PMC in 14.5% of the studied subjects. It was found in 8.3% (7/84) of the cases with an AMO, with 11.9% (5/42) being on the right side, and 2.9% (2/84) bilateral. In patients without an AMO, PMC was found in 4.9% (35/716) of the cases, with 3.6% (13/358) on the right side, 1.4% (5/358) on the left side, and 2.4% (17/716) bilateral. The incidence of PMC increased by 20% in the presence of an AMO (OR = 1.20) (►Table 1).

The Relationship between AMO and MRC

We identified MRC in 8.5% of the studied subjects. In 2.4% (2/84) of the cases, MRC was found along with an AMO. It was found in 5.6% (40/716) of the cases without an AMO, with 4.75% (17/358) being on the right side, 1.4% (5/358) on the left side, and 2.5% (18/716) bilateral. The incidence of MRC decreased by 50% in the presence of an AMO (OR = 0.50) (►Table 1).

The Relationship between AMO and MS

We identified MS in 49.8% of the studied subjects. It was found in 28.6% (24/84) of the cases with an AMO, 23.8%




Fig. 5 Axial (A) and Coronal (B) scans of mucus retention cyst. C –mucosal thickening in right side; D –maxillary sinusitis.

(10/42) of which were on the right side, 7.1% (3/42) on the left side, and 13.1% (11/84) bilateral. In patients without an AMO, MS was found in 2.5% (18/716) of the cases, with 2.2% (8/358) being on the right side, 0.5% (2/358) on the left side, and 1.1% (8/716) bilateral. The incidence of MS increased by 39% in the presence of an AMO (OR = 1.39) (► **Table 1**).

The Relationship between AMO and MT

We diagnosed 24% of the studied subjects with MT. It was found in 11.9% (10/84) of the cases with an AMO, 14.3% (6/42) of which were on the right side and 4.7% (4/84) bilateral. In patients without an AMO, MT was found in 4.5% (32/716) of the cases, with 3.3% (12/358) being on the right side, 1.4% (5/358) on the left side and 2.1% (15/716) bilateral. Therefore, AMO did not increase nor decrease MT (OR = 0.98) (► **Table 1**).

Discussion

Although the literature contains several studies of the incidence of AMO, to the best of our knowledge, only one other article⁸ discusses the presence of an AMO and anatomical variations in adjacent structures and sinus diseases. Thus, the present study aimed to investigate the relationship between the incidence of AMO and the many important variations in adjacent structures, along with the presence of sinusitis. In studies performed with cadavers and patients, the reported prevalence of AMO was in the range of 0 to 43%.^{18,19} It was reported as 10% by May et al.,²⁰ 15% by Kennedy and Zinteich,²¹ 18.5% by Singhal and Singhal,²² 19.1% by Yenigun et al.,⁸ 23% by Van Alyea,²³ and 43% by Schaeffer.²⁴ However, some of the current studies have shown results above this range. Yeung et al.²⁵ and Hung

Table 1 Concurrence of accessory maxillary ostium with neighboring morphologic variants

Variations	Presence/absence of right AMO N (%)	Presence/absence of left AMO N (%)	Presence/absence of bilateral AMO N (%)	Presence/absence of total AMO N (%)	OR
Agger nasi cell	12–6 (28.6–1.7)	1–4 (2.4–1.1)	13–6 (15.5–0.8)	26–16 (31–2.2)	1.77
Haller cell	1–17 (2.4–4.75)	0–5 (0–1.4)	4–15 (4.7–2.1)	5–37 (5.9–5.16)	1.20
Nasal septum deviation	10–8 (23.8–2.2)	0–5 (0–1.4)	10–9 (11.9–1.25)	20–22 (23.8–3.07)	1.32
Hypertrophy of inferior concha	10–8 (23.8–2.2)	0–5 (0–1.4)	9–10 (10.7–1.4)	19–23 (22.6–3.2)	1.61
Pneumatization of middle concha	5–13 (11.9–3.6)	0–5 (0–1.4)	2–17 (2.9–2.4)	7–35 (8.3–4.9)	1.20
Mucus retention cysts	1–17 (2.4–4.75)	0–5 (0–1.4)	1–18 (1.2–2.5)	2–40 (2.4–5.6)	0.50
Maxillary sinusitis	10–8 (23.8–2.2)	3–2 (7.1–0.5)	11–8 (13.1–1.1)	24–18 (28.6–2.5)	1.39
Mucosal thickening	6–12 (14.3–3.3)	0–5 (0–1.4)	4–15 (4.7–2.1)	10–32 (11.9–4.5)	0.98

Abbreviations: AMO, accessory maxillary ostium; OR, odds ratio.

Note: In patients with AMO, the first number in the table represents the number of patients with corresponding left column variation, and the second number represents the number of patients without variation.

et al.²⁶ reported the incidence of AMO as 45.5% and 47.2%, respectively. In the present study, the incidence (10.5%) was found to be slightly lower, although within the range reported in previous studies.

Avsever et al.²⁷ reported the incidence of ANC, HC, NSD and PMC as 2.9%, 3.2%, 13.2% and 13.7%, respectively. The present study found higher results. Arslan et al.²⁸ reported the incidence of AMO, MRC and NSD as 30%, 18.2% and 58.7%, respectively. The parameters reported in our study were shown to be lower. In the study of Yenigun et al.,⁸ the incidence of right, left, and bilateral AMO was 7.2%, 3.7%, and 8.2%, respectively. Kumar et al.¹⁸ reported that AMO incidence was twice as high on the right side compared with the left. Similarly, in the present study, an AMO incidence was found in 4.5% of the patients on the right side, in 1.25% of the patients on the left side, and in 4.75% of the patients bilaterally.

The incidence of ANC was reported as 51.9% by Özdemir et al.,²⁹ 40% by Orhan and Saylam,³⁰ 62.8% by Yenigun et al.⁸ and 81.8% by Liu et al.³¹ In the present study, ANC was diagnosed in 49.3% of the studied subjects. In 31% of the cases, it was diagnosed in the presence of AMO, with a 77% increase. By contrast, Yenigun et al.⁸ reported that there was no statistical significance in the simultaneous presence of AMO and ANC.

The incidence of HC has been reported to vary from 2 to 45% in the literature.³² In the present study, the incidence was found to be 10.5%, which places it within the reported limits. In 5.9% of the cases, it was found along with AMO, with a 20% increase. By contrast, Yenigun et al.⁸ reported that there was no statistical significance for the simultaneous presence of AMO and HC.

Some studies have reported NSD in from 20 to 31% of the community, and also found that severe deviation predisposed the population to rhinosinusitis.^{33–35} In Turkey, some studies reported the incidence of NSD as 39%³⁶ in adults and

34.9%³⁷ in children. Yenigun et al.⁸ reported the incidence of NSD as 47.7%. In present study, the incidence was 41.5%, similar to other studies in the literature. However, in the studies by de Oliveira et al.,³⁸ Stallman et al.³⁹ and Clark et al.,⁴⁰ a higher incidence (60.3%, 65% and 76%, respectively) was reported. In our study, NSD was found along with AMO in 23.8% of the cases. In contrast to the present study, Yenigun et al.⁸ reported that there was no statistical significance for the simultaneous presence of AMO and NSD.

The incidence of HIC was reported as 37.4% by Yenigun et al.,⁸ 72% by Clark et al.⁴⁰ and 6.83% by Cury et al.⁴¹ The present study found that HIC was diagnosed in 35% of the studied subjects, with 22.6% being diagnosed along with AMO. The low incidence in the study of Cury et al.⁴¹ may be due to the use of panoramic radiography as the radiological method. Yenigun et al.⁸ reported that the simultaneous presence of AMO and HIC was statistically significant on the left side but not significant on the right side. In this study, there was a 61% increasing incidence of HIC in the presence of AMO.

The incidence of PMC was reported in the range between 13 and 72.2% in the literature.^{42–44} Stallman et al.³⁹ and Yenigun et al.⁸ reported the incidence of PMC as 44% and 44.9%, respectively. In the present study, PMC was diagnosed in 14.5% of the studied subjects, 8.3% of whom were also diagnosed with AMO, with an increase of 20%. By contrast, Yenigun et al.⁸ reported that there was no statistical significance for the simultaneous presence of AMO and PMC.

The diagnosis of MRC in the maxillary sinus is frequent, and radiological studies reported its incidence as 9 to 22 % in the general population, similar to our findings of 8.5%.⁴⁵ Yenigun et al.⁸ reported that the presence of AMO was associated with an approximate 3-fold increase in the incidence of MRC. However, in this study, a 50% decreasing incidence of MRC in the presence of AMO was reported.

The incidence of MT was found to be 25% by Yenigun et al.,⁸ 38.1% by Ritter et al.,⁴⁶ 35.1% by Raghav et al.,⁴⁷ 21.25%

by Drumond et al.,⁴⁸ and 25.1% by Gracco et al.⁴⁹ Compatible with these studies, our study indicated that MT was diagnosed in 24% of the studied subjects, and in 14.3% it was found along with AMO. MT is formed in paranasal sinuses after infection out of maxillary sinuses, whereas the development of MRC is very rarely seen.^{50,51} In the present study, the incidence of MT was found to be higher than that of MRC, possibly for the same reason. Yenigun et al.⁸ reported that the presence of AMO was associated with a nearly 2-fold increase in the incidence of MT. By contrast, in our study, the presence of AMO did not increase nor decrease incidence of MT.

The most frequent disease of the paranasal sinuses is MS, and the incidence of odontogenic MS ranges from 10 to 40% of all MS.⁵² Drumond et al.⁴⁸ found the lower incidence of chronic sinusitis and chronic odontogenic sinusitis to be 7.48% and 2.29%, respectively. Yenigun et al.⁸ reported the incidence of MS as 14.5%. A higher incidence of 49.8% was found in our study, 28.6% of which it was diagnosed along with AMO. Yenigun et al.⁸ reported that the presence of AMO was associated with a nearly 2-fold increase in the incidence of MS. Bani-Ata et al.⁵³ reported that the presence of AMO can contribute to the occurrence of MS. Similarly, in our study, a 39% increasing incidence of MS in the presence of AMO was observed.

The differences in results may be due to the large number of parameters in our study, their effects on one another, and therefore the extensive literature review involving evaluation of studies conducted in different populations, using different numbers of patients and different radiological modalities.

Vital structures such as the skull base, optic nerves, internal carotid arteries, and orbits lie near the paranasal sinuses. Some anatomic variants increase the risk of injury in these structures, and a clear understanding of the patient's individual paranasal sinus anatomy obtained by CT examination is very important for safe endoscopic sinus surgery.⁵⁴ In most cases, CT is accepted as a gold standard modality to diagnose the sinus diseases, as multiple contiguous thin sections in axial, sagittal and coronal planes can be obtained. Bone and soft tissue observation can also be performed.⁵⁵

It should also be noted that MS can have life-threatening complications, such as orbital, intracranial, or combinations thereof. Intracranial complications of sinusitis are diseases that require an emergency approach, early diagnosis, and intensive treatment.^{56,57}

Conclusion

This study involved sinus diseases without polyps. The results showed that the incidence of all parameters, except for MRC and MT, increased in the presence of AMO. Thus, in cases with AMO, it is vital that the clinician be aware of maxillary sinus variations to avoid complications during maxillary sinus surgery. Although the literature contains several studies on the anatomy and variations in the maxillary sinus, the present study is, to the best of our knowledge, one of the most detailed studies of the presence of AMO and anatomical variations in adjacent structures, and their rela-

tion to sinus diseases. Hence, we believe that our study will be a guide for future research.

Conflict of Interests

The authors have no conflict of interests to declare.

References

- 1 Stammberger H. Functional endoscopic sinus surgery. The Messerklinger technique. Philadelphia: BC Decker 1990;247:63–7
- 2 Hollinshed WH, Rosse C. Text book of Anatomy. 4th edn. Herper and Row Philadelphia 1985:976–85
- 3 Na Y, Kim K, Kim SK, Chung SK. The quantitative effect of an accessory ostium on ventilation of the maxillary sinus. *Respir Physiol Neurobiol* 2012;181(01):62–73. Doi: 10.1016/j.resp.2012.01.013
- 4 Joe JK, Ho SY, Yanagisawa E. Documentation of variations in sinonasal anatomy by intraoperative nasal endoscopy. *Laryngoscope* 2000;110(2 Pt 1):229–235. Doi: 10.1097/00005537-200002010-00008
- 5 Jones NS. CT of the paranasal sinuses: a review of the correlation with clinical, surgical and histopathological findings. *Clin Otolaryngol Allied Sci* 2002;27(01):11–17. Doi: 10.1046/j.0307-7772.2001.00525.x
- 6 Matthews BL, Burke AJ. Recirculation of mucus via accessory ostia causing chronic maxillary sinus disease. *Otolaryngol Head Neck Surg* 1997;117(04):422–423. Doi: 10.1016/S0194-5998(97)70139-6
- 7 Parsons DS, Wald ER. Otitis media and sinusitis: similar diseases. *Otolaryngol Clin North Am* 1996;29(01):11–25
- 8 Yenigun A, Fazliogullari Z, Gun C, Uysal II, Nayman A, Karabulut AK. The effect of the presence of the accessory maxillary ostium on the maxillary sinus. *Eur Arch Otorhinolaryngol* 2016;273(12):4315–4319. Doi: 10.1007/s00405-016-4129-8
- 9 Yegin Y, Çelik M, Şimşek BM, Olgun B, Canpolat S, Kayhan FT. Relationship Between the Presence of Agger Nasi Cells and the Development of Frontal Sinusitis: a Computer-Assisted Anatomic Study. *Bezmialem Science* 2017;5:112–115. Doi: 10.14235/bs.2017.1087
- 10 Mathew R, Omami G, Hand A, Fellows D, Lurie A. Cone beam CT analysis of Haller cells: prevalence and clinical significance. *Dentomaxillofac Radiol* 2013;42(09):20130055. Doi: 10.1259/dmfr.20130055
- 11 Vig KW. Nasal obstruction and facial growth: the strength of evidence for clinical assumptions. *Am J Orthod Dentofacial Orthop* 1998;113(06):603–611. Doi: 10.1016/s0889-5406(98)70219-7
- 12 Hatipoğlu HG, Cetin MA, Yüksel E. Concha bullosa types: their relationship with sinusitis, ostiomeatal and frontal recess disease. *Diagn Interv Radiol* 2005;11(03):145–149
- 13 Chiesa Estomba C, Rivera Schmitz T, Ossa Echeverri CC, Betances Reinoso FA, Osorio Velasquez A, Santidrian Hidalgo C. Compensatory hypertrophy of the contralateral inferior turbinate in patients with unilateral nasal septal deviation. A computed tomography study. *Otolaryngol Pol* 2015;69(02):14–20. Doi: 10.5604/00306657.1149568
- 14 Fehrenbach MJ, Herring SW. *Illustrated Anatomy of the Head and Neck*. Elsevier Health Sciences; 2012 page 67
- 15 Bhattacharyya N. Do maxillary sinus retention cysts reflect obstructive sinus phenomena? *Arch Otolaryngol Head Neck Surg* 2000;126(11):1369–1371. Doi: 10.1001/archotol.126.11.1369
- 16 Kolo ES. The role of plain radiographs in the diagnosis of chronic maxillary rhinosinusitis in adults. *Afr Health Sci* 2012;12(04):459–463. Doi: 10.4314/ahs.v12i4.10
- 17 Ren S, Zhao H, Liu J, Wang Q, Pan Y. Significance of maxillary sinus mucosal thickening in patients with periodontal disease. *Int Dent J* 2015;65(06):303–310. Doi: 10.1111/idj.12186

- 18 Kumar H, Choudhry R, Kakar S. Accessory Maxillary Ostia: Topography and Clinical Application. *J Anat Soc India* 2001;50(01):3–5
- 19 Jog M, McGarry GW. How frequent are accessory sinus ostia? *J Laryngol Otol* 2003;117(04):270–272. Doi: 10.1258/00222150360600869
- 20 May M, Sobol SM, Korzec K. The location of the maxillary os and its importance to the endoscopic sinus surgeon. *Laryngoscope* 1990;100(10 Pt 1):1037–1042. Doi: 10.1288/00005537-199010000-00002
- 21 Kennedy DW, Zintech J. *Otolaryngology, Head and Neck, Vol III*. W.B. Saunders Company; 1991:1861–71
- 22 Singhal MD, Singhal DM. Anatomy of accessory maxillary sinus ostium with clinical application. *Int J Med Sci Public Health* 2014;3:327–329. Doi: 10.5455/ijmsph.2013.301220131
- 23 Van Alyea OE. The Ostium Maxillare anatomic study of its surgical accessibility. *Arch Otolaryngol* 1936;24(05):553–569
- 24 Schaeffer JP. *The Nose, Paranasal Sinuses, Nasolacrimal Passages, and Olfactory Organ in Man*. Philadelphia: P. Blakiston's Son; 1920
- 25 Yeung AWK, Colsoul N, Montalvao C, Hung K, Jacobs R, Bornstein MM. Visibility, location, and morphology of the primary maxillary sinus ostium and presence of accessory ostia: a retrospective analysis using cone beam computed tomography (CBCT). *Clin Oral Investig* 2019;23(11):3977–3986. Doi: 10.1007/s00784-019-02829-9
- 26 Hung K, Montalvao C, Yeung AWK, Li G, Bornstein MM. Frequency, location, and morphology of accessory maxillary sinus ostia: a retrospective study using cone beam computed tomography (CBCT). *Surg Radiol Anat* 2020;42(02):219–228. Doi: 10.1007/s00276-019-02308-6
- 27 Avsever H, Gunduz K, Karakoç O, Akyol M, Orhan K. Incidental findings on cone-beam computed tomographic images: paranasal sinus findings and nasal septum variations. *Oral Radiol* 2018;34(01):40–48. Doi: 10.1007/s11282-017-0283-y
- 28 Arslan İB, Uluyol S, Demirhan E, Kozcu SH, Pekçevik Y, Çukurova İ. Paranasal Sinus Anatomic Variations Accompanying Maxillary Sinus Retention Cysts: A Radiological Analysis. *Turk Arch Otorhinolaryngol* 2017;55(04):162–165. Doi: 10.5152/tao.2017.2759
- 29 Özdemir A, Arslan S. Incidence of agger nasi and frontal cells and their relation to frontal sinusitis in a Turkish population: a CT study. *Anatomy* 2018;12(02):71–75. Doi: 10.2399/ana.18.050
- 30 Orhan M, Saylam CY. Anatomical analysis of the prevalence of agger nasi cell in the Turkish population. *Kulak Burun Bogaz Ihtis Derg* 2009;19(02):82–86
- 31 Liu SC, Wang CH, Wang HW. Prevalence of the uncinate process, agger nasi cell and their relationship in a Taiwanese population. *Rhinology* 2010;48(02):239–244. Doi: 10.4193/Rhin09.118
- 32 Stackpole SA, Edelstein DR. The anatomic relevance of the Haller cell in sinusitis. *Am J Rhinol* 1997;11(03):219–223. Doi: 10.2500/105065897781751910
- 33 Shin HS. Clinical significance of unilateral sinusitis. *J Korean Med Sci* 1986;1(01):69–74. Doi: 10.3346/jkms.1986.1.1.69
- 34 Lebowitz RA, Brunner E, Jacobs JB. The agger nasi cell: radiological evaluation and endoscopic management in chronic frontal sinusitis. *Operative techniques. Otolaryngol Head Neck Surg* 1995;6:171–175
- 35 Wanamaker HH. Role of Haller's cell in headache and sinus disease: a case report. *Otolaryngol Head Neck Surg* 1996;114(02):324–327. Doi: 10.1016/s0194-5998(96)70196-1
- 36 Özkırış M, Mutlu C. Kulak burun boğaz polikliniğine başvuran hastalarda nazal septum deviasyonu sıklığı. *J Kartal TR* 2010;XXI(02):72–76
- 37 Yıldırım I, Okur E. The prevalence of nasal septal deviation in children from Kahramanmaraş, Turkey. *Int J Pediatr Otorhinolaryngol* 2003;67(11):1203–1206. Doi: 10.1016/j.ijporl.2003.07.005
- 38 Oliveira AKP, Elias E Jr, Santos LV, Bettega SG, Mocellin M. Prevalence of deviated nasal septum in Curitiba, Brasil. *Int Arch Otorhinolaryngol* 2005;9(04):288–292
- 39 Stallman JS, Lobo JN, Som PM. The incidence of concha bullosa and its relationship to nasal septal deviation and paranasal sinus disease. *AJNR Am J Neuroradiol* 2004;25(09):1613–1618
- 40 Clark DW, Del Signore AG, Raithatha R, Senior BA. Nasal airway obstruction: Prevalence and anatomic contributors. *Ear Nose Throat J* 2018;97(06):173–176. Doi: 10.1177/014556131809700615
- 41 Cury R, Gioseffi C, Andrade E, Cury M, Cury SE. The incidence of inferior turbinate hypertrophy in a Brazilian population. *Webmed central. Rhinology* 2013;4(06):WMC004276
- 42 Aktas D, Kalcioğlu MT, Kutlu R, Ozturan O, Oncel S. The relationship between the concha bullosa, nasal septal deviation and sinusitis. *Rhinology* 2003;41(02):103–106
- 43 Arslan H, Aydinlioğlu A, Bozkurt M, Egeli E. Anatomic variations of the paranasal sinuses: CT examination for endoscopic sinus surgery. *Auris Nasus Larynx* 1999;26(01):39–48. Doi: 10.1016/s0385-8146(98)00024-8
- 44 Bolger WE, Butzin CA, Parsons DS. Paranasal sinus bony anatomic variations and mucosal abnormalities: CT analysis for endoscopic sinus surgery. *Laryngoscope* 1991;101(1 Pt 1):56–64. Doi: 10.1288/00005537-199101000-00010
- 45 Albu S. Symptomatic maxillary sinus retention cysts: should they be removed? *Laryngoscope* 2010;120(09):1904–1909. Doi: 10.1002/lary.21040
- 46 Ritter L, Lutz J, Neugebauer J, et al. Prevalence of pathologic findings in the maxillary sinus in cone-beam computerized tomography. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;111(05):634–640. Doi: 10.1016/j.tripleo.2010.12.007
- 47 Raghav M, Karjodkar FR, Sontakke S, Sansare K. Prevalence of incidental maxillary sinus pathologies in dental patients on cone-beam computed tomographic images. *Contemp Clin Dent* 2014;5(03):361–365. Doi: 10.4103/0976-237X.137949
- 48 Drumond JPN, Allegro BB, Novo NF, de Miranda SL, Sendyk WR. Evaluation of the prevalence of maxillary sinuses abnormalities through spiral computed tomography (CT). *Int Arch Otorhinolaryngol* 2017;21(02):126–133. Doi: 10.1055/s-0036-1593834
- 49 Gracco A, Incerti Parenti S, Ioele C, Alessandri Bonetti G, Stellini E. Prevalence of incidental maxillary sinus findings in Italian orthodontic patients: a retrospective cone-beam computed tomography study. *Korean J Orthod* 2012;42(06):329–334. Doi: 10.4041/kjod.2012.42.6.329
- 50 Wang JH, Jang YJ, Lee BJ. Natural course of retention cysts of the maxillary sinus: long-term follow-up results. *Laryngoscope* 2007;117(02):341–344
- 51 Hong SL, Cho KS, Roh HJ. Maxillary sinus retention cysts protruding into the inferior meatus. *Clin Exp Otorhinolaryngol* 2014;7(03):226–228. Doi: 10.3342/ceo.2014.7.3.226
- 52 Mehra P, Murad H. Maxillary sinus disease of odontogenic origin. *Otolaryngol Clin North Am* 2004;37(02):347–364. Doi: 10.1016/S0030-6665(03)00171-3
- 53 Bani-Ata M, Aleshawi A, Khatatbeh A, et al. Accessory maxillary ostia: prevalence of an anatomical variant and association with chronic sinusitis. *Int J Gen Med* 2020;13:163–168. Doi: 10.2147/IJGM.S253569
- 54 Nouraei SA, Elisay AR, Dimarco A, et al. Variations in paranasal sinus anatomy: implications for the pathophysiology of chronic rhinosinusitis and safety of endoscopic sinus surgery. *J Otolaryngol Head Neck Surg* 2009;38(01):32–37
- 55 Campbell PD Jr, Zinreich SJR, Aygun N. Imaging of the paranasal sinuses and in-office CT. *Otolaryngol Clin North Am* 2009;42(05):753–764, vii
- 56 Clayman GL, Adams GL, Paugh DR, Koopmann CF Jr. Intracranial complications of paranasal sinusitis: a combined institutional review. *Laryngoscope* 1991;101(03):234–239. Doi: 10.1288/00005537-199103000-00003
- 57 Giannoni C, Sulek M, Friedman EM. Intracranial complications of sinusitis: a pediatric series. *Am J Rhinol* 1998;12(03):173–178. Doi: 10.2500/105065898781390127