

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/274274026>

Assessment of aortic elasticity and its relation with left ventricular filling and diastolic parameters by transthoracic and transesophageal echocardiography

Article · January 2013

DOI: 10.5505/abantmedj.2013.08370

CITATIONS

0

READS

12

6 authors, including:



Ibrahim Halil Tanboga

Ataturk University

189 PUBLICATIONS 1,459 CITATIONS

[SEE PROFILE](#)



Mehmet Fatih Karakas

Mustafa Kemal University

69 PUBLICATIONS 552 CITATIONS

[SEE PROFILE](#)



Eyup Buyukkaya

Mustafa Kemal University

42 PUBLICATIONS 508 CITATIONS

[SEE PROFILE](#)

Assessment of Aortic Elasticity and Its Relation with Left Ventricular Filling and Diastolic Parameters by Transthoracic and Transesophageal Echocardiography

Aortik Elastisite ve Aortik Elastisitenin Sol Ventrikül Dolum ve Diyastolik Parametreleri ile İlişkisinin Transtorasik ve Transözofajiyal Ekokardiyografi Değerlendirilmesi

Mustafa Kurt¹, İbrahim Halil Tanboga², Mehmet Fatih Karakaş¹, Eyüp Büyükkaya¹, Adnan Burak Akcay¹, Nihat Sen¹

¹Mustafa Kemal Üniversitesi, Tıp Fakültesi, Kardiyoloji Ana Bilim Dalı, Hatay

²Atatürk Üniversitesi, Tıp Fakültesi, Kardiyoloji Ana Bilim Dalı, Erzurum

Abstract

Background: There are limited data comparing aortic dimensions and elastic properties between transthoracic echocardiography (TTE) and transesophageal echocardiography (TEE). Therefore, we aimed to investigate aortic dimensions and indices of aortic elastic properties as well as their relation with left ventricular (LV) filling and diastolic parameters obtained from TTE and TEE.

Method: Forty two patients were included in the study. Aortic dimensions and related elasticity parameters, LV filling and diastolic parameters, left atrial volume index (LAVi), LV mass index were calculated from TTE and TEE recordings in all patients. The correlation analyses were performed for aortic elasticity indices and LV filling and diastolic parameters obtained from TTE and TEE.

Results: There were no significant differences in aortic dimensions and elasticity indexes obtained from TTE and TEE. The correlation analysis revealed that while age E/A, E/E' and LAVi were significantly correlated with aortic elasticity indexes obtained from both TTE and TEE, LV mass index was significantly correlated with only TEE aortic elasticity indices.

Conclusion: When compared to TTE, aortic elastic properties obtained from TEE were more closely related with LV filling and diastolic parameters. In addition, the indexes of aortic function were comparable between TEE and TTE with good intra and interobserver variability.

Keywords: aortic dimensions, transthoracic echocardiography and transesophageal echocardiography.

Özet

Amaç: Aort boyutları ve elastik özelliklerini transtorasik ekokardiyografi (TTE) ve transözofajiyal ekokardiyografi (TEE) ile karşılaştıran sınırlı veri vardır. Bu nedenle, TTE ve TEE elde edilen aort boyutları ve aortik elastik indekslerinin karşılaştırmanın yanı sıra bunların sol ventrikül (LV) dolum ve diyastolik parametreleri ile ilişkisini araştırmayı amaçladık.

Yöntem: Çalışmaya kırk iki hasta dahil edildi. Tüm hastaların TTE ve TEE kayıtlarından aort boyutları ve ilgili elastikiyet parametreleri, sol ventrikül dolum ve diyastolik parametreleri, sol atriyal volüm indeksi (LAVi), sol ventrikül kütle indeksi hesaplandı. TTE ve TEE elde edilen aort elastikiyeti indeksleri ve sol ventrikül dolum ve diyastolik parametreleri için korelasyon analizi yapıldı.

Bulgular: TTE ve TEE ile elde edilen aort boyutları ve elastikiyeti indeksleri arasında anlamlı bir fark yoktu. Korelasyon analizinde yaş, E / A, E / E' ve LAVi hem TTE hem de TEE ile elde edilen aort elastikiyeti indeksleri ile anlamlı derecede ilişkili iken, sol ventrikül kütle indeksi sadece TEE ile elde edilen aort elastikiyeti indeksleri ile ilişkili saptandı.

Sonuç: TTE ile karşılaştırıldığında, TEE elde edilen aortik elastik özellikleri sol ventrikül dolum ve diyastolik parametreleri ile daha yakından ilişkili bulunmuştur. Buna ek olarak, TEE ve TTE ile elde edilen aort fonksiyon indeksleri iyi gözlemci içi ve gözlemciler arası değişkenliği ile benzerdir.

Anahtar Kelimeler: aort boyutları, transtorasik ve transözofajiyal ekokardiyografi.

Introduction

Impaired aortic elasticity may be cause the increase of afterload. Increasing the afterload can be also increased left ventricular filling pressure and impaired of LV systolic and diastolic function(1-5). Impaired aortic elasticity may contribute to the development of left ventricular diastolic and systolic dysfunction due to increased pulse pressure and left ventricular afterload which results in left ventricular hypertrophy and subendocardial ischemia (6). To quantify the aortic elasticity, many techniques can be used such as transthoracic echocardiography (TTE) or

transesophageal echocardiography (TEE), magnetic resonance imaging, and computed tomography (CT)(7-9). However, there are very limited studies related to assessment of aortic elasticity on TEE (10-16). Moreover, in majority of these studies, the elastic properties of aorta were measured on descending aorta in contrast to TTE studies and there were only a few data regarding to elasticity indices of ascending aortic by TEE. In addition, there is no data comparing aortic elastic properties on TTE and TEE and their relation with LV diastolic parameters. The purpose of this study was to compare indices of aortic elastic properties obtained from patients undergoing simultaneous TTE and TEE and to determine their relation with LV diastolic parameters.

İletişim Bilgisi / Correspondence

Yard. Doç. Dr. Mustafa Kurt, Mustafa Kemal Üniversitesi; Araştırma Ve Uygulama Hastanesi, Serinyol, Hatay E-posta: drmustafakurt@yahoo.com

Geliş tarihi / Received: 27.11.2012; Kabul tarihi / Accepted: 14.12.2012 Çıkar Çatışması / Conflict Of Interest: Yok /None

Material and Methods

Study Patients

Forty-two patients underwent a clinically indicated study. The patients with atrial fibrillation, previous ischemic heart disease, LV hypertrophy, left bundle branch block, pericardial disease, poor image quality and inability to give consent were excluded from the study. TTE and TEE was indicated for 31 patients included in the study for the evaluation of patent foramen ovale, 11 patients for the evaluation of bicuspid aortic valve. The study protocol was approved by the institutional review board, and patients provided a written informed consent.

Transthoracic Echocardiography

Transthoracic echocardiography, with a 2.5 MHz phased-array transducer, was performed using a GE Vivid 7 system (GE Vingmed Ultrasound, Horten, Norway). Three consecutive measurements were obtained and the results were averaged. Recordings were taken in the left lateral decubitus position. The Doppler signals were also recorded at a speed of 100 mm/s and the M-mode traces were recorded at a speed of 50 mm/s (17). Simultaneous electrocardiographic recordings were also taken.

Ascending aorta diameter was measured on the M-mode tracing at 3 cm above the aortic valve. The systolic diameter was measured at the maximal anterior motion of the aorta, whereas the diastolic diameter was measured at the peak of the QRS complex on the simultaneously recorded electrocardiogram. By the biplane Simpson's method LV ejection fraction (LVEF) was calculated. According to the recommendations of the American Society of Echocardiography LV filling and diastolic parameters such as left atrial volume index (LAVi), E, A, Em waves and Deceleration time (DecT) were calculated (18). Besides, LV mass was calculated according to previously suggestions (19).

Transesophageal Echocardiography

TEE, with a 5.0-MHz transesophageal probe, was performed by using a GE Vivid 7 system (GE

Vingmed Ultrasound AS, Horten, Norway). TEE was performed after 4 hours fasting period just after transthoracic echocardiography. For posterior pharyngeal anesthesia ten percent lidocain spray was used. The procedure was performed with continuous monitoring of blood pressure, heart rate, one lead ECG and saturation. Ascending aortic elastic parameters was calculated from midesophageal long axis view with approximately 120°. Diameter of the ascending aorta was measured on the M-mode tracing at 3 cm above of the aortic valve. The systolic diameter was measured as previously mentioned at the maximal anterior motion of the aorta, whereas the diastolic diameter was measured at the peak of the QRS complex on the simultaneously recorded electrocardiogram

Aortic Elasticity parameters

For aortic elasticity, following formulas were used (20): aortic strain (%) = [(aortic systolic diameter – diastolic diameter) × 100] / aortic diastolic diameter, aortic stiffness index = $\ln(\text{systolic pressure} / \text{diastolic pressure}) / \text{aortic strain}$, aortic distensibility ($\text{cm}^2 \cdot \text{dyn}^{-1} \cdot 10^{-6}$) = $(2 \times \text{aortic strain}) / (\text{systolic pressure} - \text{diastolic pressure})$ and aortic elastic modulus ($\text{dyn} \cdot \text{cm}^{-2} \cdot 10^6$) = $(\text{systolic pressure} - \text{diastolic pressure}) / \text{aortic strain}$.

Reproducibility

The intra- and inter-observer variability of aortic dimension measurements were determined for TTE and TEE in all patients. For intraobserver variability, the same observer repeated the measurements after two weeks, whereas interobserver variability was assessed from the measurements by two independent observers.

Statistics

Continuous variables are expressed as mean ± SD. The level of significance was accepted as $p < 0.05$. For the normality test of all variables the Kolmogorov-Smirnov test was used. For compare TTE and TEE parameters, independent Student t test or the Mann-Whitney U test was used. Correlation between variables was tested by Pearson or Spearman correlation tests. Interobserver and intraobserver variability assessed by Bland-Altman analysis and intraclass correlation coefficient.



cient (ICC). An ICC greater than 0.75 represents excellent reproducibility, whereas values between 0.4 and 0.75 represent fair to good reproducibility (21).

Results

The study population consisted of 42 patients (mean age 36.2 ± 11.9 years and 48% male).. Patients' mean body mass index were 25.5 ± 5.1 , mean LV-EF were 64 ± 3.7 , mean LV-Mass index were 105 ± 18 and mean LAVi were 25 ± 10.5 . The other characteristics were given in table 1 and 2. There were no significant differences between the TTE and TEE in terms of systolic and diastolic blood pressure and heart rate ($p=0.32$, $p=0.42$ and $p=0.58$, respectively).

Table 1. Basic characteristics of the patients.

Age – years	36.2±11.9
Male (%)	48
Smoking (%)	24
Hypertension (%)	12
Body mass index (kg/m ²)	25.5±5.1
Body surface Area (m ²)	1.79±0.16
Systolic blood pressure (mmHg)	116±22
Diastolic blood pressure (mmHg)	71±13
Heart rate – per minute	82±12

Table 2. Basic Echocardiographic characteristics of the patients.

Left ventricular ejection fraction (%)	64±3.7
Left ventricular Mass index (gr/m ²)	105±18
Left atrial volume index (ml/m ²)	25±10.5
E wave (m/sec)	0.75±0.19
A wave (m/sec)	0.68±0.22
E/A ratio	1.2±0.36
E' wave (m/sec)	0.11±0.03
E/E' ratio	6.7±1.7
Deceleration time (msec)	186±43

There were no significant differences between the TTE and TEE in terms of aortic elasticity parameters (Table 3). There was a moderate correlation with TTE aortic strain E/A, e/e' , LAVi and age, and a weak correlation with LV-Mass-i and DecT. TEE aortic strain showed a moderate correlation with all parameters except for DecT. TTE aortic distensibility had a moderate correlation

with E/A, e/e' , LAVi and age, and a weak correlation with LV-Mass-i and DecT. TEE aortic distensibility was shown to have a weak correlation with DecT, a moderate one with LV Mass-i, and a strong one with the other parameters.

Table 3. Comparison of aortic elasticity parameters by TTE and TEE.

Variable	TTE	TEE	p
Aortic systolic diameter (mm)	27.4±2.6	27.3±2.7	0.95
Aortic diastolic diameter (mm)	25.6±2.9	25.9±3.0	0.67
Aortic strain (%)	7.1±4.4	5.5±2.9	0.12
Aortic Distensibility (cm ² .dyn ⁻¹ .10 ⁻⁶)	3.5±3.0	2.5±1.48	0.12
Aortic Elastic modulus (dyn.cm ⁻² .10 ⁶)	8.5±4.3	11±7.3	0.14
Aortic stiffness index	9.2±4.7	11.7±6.8	0.15
Aortic Sclerosis (Grade 0 - %)	76	56	0.13

As for TTE aortic stiffness, a moderate correlation was found with e/e' and LAVi, and a weak correlation with the other parameters. TEE aortic stiffness had a strong correlation with age, a weak correlation with DecT and a moderate correlation with the other parameters. TTE-aortic elastic modulus had a moderate correlation with E/A, e/e' , LAVi and age and a weak correlation with LV-Mass-i ve DecT. TEE-aortic elastic modulus was strongly with age, weakly correlated with DecT and moderately with the other parameters (Table 4). Inter and intraobserver agreement for aortic dimensions obtained from TTE and TEE were good (Table-5).

Discussion

The results of present study demonstrated that indexes of aortic function such as aortic strain, distensibility, stiffness and elastic modulus, obtained by TEE are more correlated LV filling parameters E/A, E/E', LAVi, LV Mass-i than obtained by TTE. To the best of our knowledge, this is the first study that indexes of aortic function by measured TEE as a simple and reliable method for prediction of diastolic dysfunction in people with a normal LVEF. In our study, indexes of aortic function such as aortic stiffness were associated with a prolonged DecT, increased E/A ratio and E/Em ratio suggesting a link between increased aortic stiffness and impaired myocardial

Table 4. Correlation coefficients between aortic elasticity parameters and left ventricular filling pressure-diastolic parameters and age.

Aortic elasticity parameters								
	Strain		Distensibility		Stiffness		Elastic modulus	
	TTE	TEE	TTE	TEE	TTE	TEE	TTE	TEE
E/A	0.43*	0.47*	0.37*	0.50*	-0.19	-0.40*	-0.28*	-0.43*
E/E'	-0.30*	-0.47*	-0.34*	-0.65*	0.28*	0.37*	0.43*	0.45*
DecT	-0.15	-0.19	-0.10	-0.17	0.14	0.21	0.09	0.17
LAVi	-0.41*	-0.45*	-0.43*	-0.53*	0.39*	0.29*	0.38*	0.26*
LVMass-i	-0.19	-0.25*	-0.20	-0.26*	0.14	0.30*	0.07	0.26*
Age	-0.43*	-0.53*	-0.39*	-0.60*	0.13	0.60*	0.26*	0.66*

*P value < 0.05 DecT: Deceleration time, LAVi: Left atrial volume index, LVMass-i: Left ventricular mass index

relaxation. E/A ratio and E/Em ratio are the marker of cardiac diastolic dysfunction, and the changes in E/A ratio seem to precede the changes in the deceleration time of mitral inflow in the early stage of cardiac diastolic dysfunction (18,22). Indexes of aortic function such as aortic distensibility are a more valid marker of arterial stiffness and better predictor of diastolic dysfunction (6,23).

The reflected wave travels more rapidly along the arterial tree in those with arterial stiffness, resulting in an increase in aortic systolic pressure and a decrease in aortic pressure during diastole (7). The resultant increase in afterload during LV systole and reduction in coronary perfusion during LV diastole may lead to LV hypertrophy and slowing of left ventricle relaxation (24). Clinical studies have also linked increased aortic stiffness to overt diastolic heart failure and coronary artery disease extension (7,25-27). Thus, abnormal aortic elastic properties may represent one of the mechanisms of LV diastolic dysfunction in our study population with normal LVEF.

Also, we showed that indexes of aortic function such as aortic stiffness were associated with age. Dilatation of blood vessel occurs not only normally with aging, but in addition, elastic properties of the arterial vessel wall decrease with advancing age (28,29). Lipid deposits and lipid metabolism in arterial walls accelerate elastic fiber destruction during the process of atherosclerosis, thus further reducing the distensibility of the aorta. Since the destroyed elastic fibers are replaced with collagen fibers, the stiffness of the aorta increases, thereby greatly affecting not only cardiovascular dynamics, but also cardiac function. Furthermore, the calcium content of the aorta that is observed in atherosclerosis also reduces the distensibility of the aorta (30).

In conclusion, aortic elastic properties obtained from TEE were more closely related with LV filling and diastolic parameters than aortic elastic properties obtained from TTE. In addition, the indexes of aortic function were comparable between TEE and TTE with good intra- and interobserver variability.

Table 5. Inter and intraobserver agreement for aortic dimension obtained from TTE and TEE.

Dimensions	Intraobserver agreement ICC, 95% CI	Interobserver agreement, Mean difference, 95% CI
TTE-Aortic systolic	0.96 (0.91-0.98)	-0.03 (0.31, -0.37)
TTE-Aortic diastolic	0.94 (0.87-0.97)	-0.09 (0.17, -0.35)
TEE-Aortic systolic	0.92 (0.83-0.97)	-0.07 (0.38, -0.52)
TEE-Aortic diastolic	0.90 (0.78-0.96)	0.03 (0.33, -0.27)

Limitations

The most important limitations of the present study are the low number of participants and not being used of TEE in routine clinical practice frequently. However, the implication of our study is that when compared to TTE aortic elasticity parameters, TEE aortic elasticity parameters were better reflects the LV diastolic function in patients who underwent TEE for any reason. Besides, non-invasive measurement of blood pressures from brachial artery could be accepted as a limitation, however in the clinical setting, it can-

not be feasible to have invasively measured central pressure. Moreover, assessment of E/A and

E/Em in patients with normal EF have some limitations.

KAYNAKLAR

1. Eren M, Gorgulu S, Uslu N, Celik S, Dagdeviren B, Tezel T. Relation between aortic stiffness and left ventricular diastolic function in patients with hypertension, diabetes, or both. *Heart* 2004;90:37-43.
2. Karamitsos TD, Karvounis HI, Didangelos TP, Papadopoulos CE, Kachrimanidou MK, Selvanayagam JB, Parharidis GE. Aortic elastic properties are related to left ventricular diastolic function in patients with type 1 diabetes mellitus. *Cardiology* 2008;109:99-104.
3. Mottram PM, Haluska BA, Leano R, Carlier S, Case C, Marwick TH. Relation of arterial stiffness to diastolic dysfunction in hypertensive heart disease. *Heart* 2005;91:1551-6.
4. Seyfeli E, Duru M, Saglam H, Akgul F, Kuvandik G, Kaya H, Yalcin F. Association of left ventricular diastolic function abnormalities with aortic elastic properties in asymptomatic patients with type 2 diabetes mellitus. A tissue doppler echocardiographic study. *Int J Clin Pract* 2008;62:1358-65.
5. Tsioufis C, Chatzis D, Dimitriadis K, Stougianos P, Kakavas A, Vlasseros I, Tousoulis D, Stefanadis C, Kalikazaros I. Left ventricular diastolic dysfunction is accompanied by increased aortic stiffness in the early stages of essential hypertension: a TDI approach. *J Hypertens* 2005;23:1745-50.
6. Safar ME, Levy BI, Struijker-Boudier H. Current perspectives on arterial stiffness and pulse pressure in hypertension and cardiovascular diseases. *Circulation* 2003;107:2864-9.
7. Laurent S, Cockcroft J, Van Bortel L, Boutouyrie P, Giannattasio C, Hayoz D, Pannier B, Vlachopoulos C, Wilkinson I, Struijker-Boudier H. Expert consensus document on arterial stiffness: methodological issues and clinical applications. *Eur Heart J* 2006;27:2588-605.
8. Ganten MK, Krautter U, von Tengg-Kobligk H, Bockler D, Schumacher H, Stiller W, Delorme S, Kauczor HU, Kauffmann GW, Bock M. Quantification of aortic distensibility in abdominal aortic aneurysm using ECG-gated multi-detector computed tomography. *Eur Radiol* 2008;18:966-73.
9. Stefanadis C, Stratos C, Boudoulas H, Kourouklis C, Toutouzas P. Distensibility of the ascending aorta: comparison of invasive and non-invasive techniques in healthy men and in men with coronary artery disease. *Eur Heart J* 1990;11:990-6.
10. Vitarelli A, Conde Y, Cimino E, D'Orazio S, Stellato S, Battaglia D, Padella V, Caranci F, Continanza G, Dettoni O, Capotosto L. Assessment of ascending aorta distensibility after successful coarctation repair by strain Doppler echocardiography. *J Am Soc Echocardiogr* 2008;21:729-36.
11. Honma H, Ohno T, Fujimoto H, Matsuzaki T, Murata H, Mizuno K. Evaluation of the elastic properties of the thoracic descending aorta with strain-rate measurement with transesophageal echocardiography: its correlation with the left ventricular diastolic function assessed with transthoracic echocardiography. *J Nippon Med Sch* 2010;77:145-54.
12. Kim KH, Park JC, Yoon HJ, Yoon NS, Hong YJ, Park HW, Kim JH, Ahn Y, Jeong MH, Cho JG, Kang JC. Usefulness of aortic strain analysis by velocity vector imaging as a new echocardiographic measure of arterial stiffness. *J Am Soc Echocardiogr* 2009;22:1382-8.
13. Vitarelli A, Conde Y, Cimino E, D'Angeli I, D'Orazio S, Stellato S, Padella V, Caranci F. Aortic wall mechanics in the Marfan syndrome assessed by transesophageal tissue Doppler echocardiography. *Am J Cardiol* 2006;97:571-7.
14. Pasiński TJ, Binkley PF, Pearson AC. Evaluation of aortic distensibility with transesophageal echocardiography. *Am Heart J* 1992;123:1288-92.
15. Petrini J, Yousry M, Rickenlund A, Liska J, Hamsten A, Eriksson P, Franco-Cereceda A, Caidahl K, Eriksson MJ. The feasibility of velocity vector imaging by transesophageal echocardiography for assessment of elastic properties of the descending aorta in aortic valve disease. *J Am Soc Echocardiogr* 2010;23:985-92.
16. Sako H, Miura S, Kumagai K, Saku K. Associations between augmentation index and severity of atheroma or aortic stiffness of the descending thoracic aorta by transesophageal echocardiography. *Circ J* 2009;73:1151-6.
17. Kurt M, Karakas MF, İpek G, Arslantaş U, Bilen E, Yüksel İÖ, Yaşar AS, Bilge M. Assessment of right ventricular function in patients with rheumatic mild mitral involvement by tissue Doppler Echocardiography. *Abant Med J*, 2012;1:39-44.
18. Nagueh SF, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA, Waggoner AD, Flachskampf FA, Pellikka PA, Evangelista A. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *J Am Soc Echocardiogr* 2009;22:107-33.
19. Foppa M, Duncan BB, Rohde LE. Echocardiography-based left ventricular mass estimation. How should we define hypertrophy? *Cardiovasc Ultrasound* 2005;3:17.
20. Ulusoy RE YM, Baysan O, Kırılmaz A, Kılıçaslan F, Cebeci BS. Ventricular functions, aortic elastic properties, and endothelial functions in patients with hypertensive response to treadmill exercise testing. *Turk J Med Sci* 2011;41:781-787.
21. Tanboga IH, Ekinci M, Isik T, Kurt M, Kaya A, Sevimli S. Reproducibility of Syntax Score: From Core Lab to Real World. *J Interv Cardiol* 2011.
22. Redfield MM, Jacobsen SJ, Burnett JC, Jr., Mahoney DW, Bailey KR, Rodeheffer RJ. Burden of systolic and diastolic ventricular dysfunction in the community: appreciating the scope of the heart failure epidemic. *JAMA* 2003;289:194-202.
23. Wang CP, Hung WC, Yu TH, Hsu HL, Chen YH, Chiu CA, Lu LF, Chung FM, Cheng YA, Lee YJ. Brachial-ankle pulse wave velocity as an early indicator of left ven-



- tricular diastolic function among hypertensive subjects. *Clin Exp Hypertens* 2009;31:31-43.
24. Roman MJ, Ganau A, Saba PS, Pini R, Pickering TG, Devereux RB. Impact of arterial stiffening on left ventricular structure. *Hypertension* 2000;36:489-94.
25. Şatıroğlu Ö BM, Bayar N, Çiçek Y, Çetin M, Bozkurt E. Relation between aortic stiffness and extension of coronary artery disease. *Turk J Med Sci*;42:417-424.
26. Weber T, O'Rourke MF, Ammer M, Kvas E, Punzengruber C, Eber B. Arterial stiffness and arterial wave reflections are associated with systolic and diastolic function in patients with normal ejection fraction. *Am J Hypertens* 2008;21:1194-202.
27. Hundley WG, Kitzman DW, Morgan TM et al. Cardiac cycle-dependent changes in aortic area and distensibility are reduced in older patients with isolated diastolic heart failure and correlate with exercise intolerance. *J Am Coll Cardiol* 2001;38:796-802.
28. Stefanadis C, Wooley CF, Bush CA, Kolibash AJ, Boudoulas H. Aortic distensibility abnormalities in coronary artery disease. *Am J Cardiol* 1987;59:1300-4.
29. Kang SM, Ha JW, Chung N, Jang KJ, Shin MS, Rim SJ, Cho SY. Assessment of elastic properties of the descending thoracic aorta by transesophageal echocardiography with acoustic quantification in patients with a stroke. *Echocardiography* 2000;17:713-20.
30. Al-Mallah MH, Nasir K, Katz R, Takasu J, Lima JA, Bluemke DA, Hundley G, Blumenthal RS, Budoff MJ. Thoracic aortic distensibility and thoracic aortic calcium (from the Multi-Ethnic Study of Atherosclerosis [MESA]). *Am J Cardiol* 2010;106:575-80.

