

Growth, yield, and fruit quality of 'Rhode Red Valencia' and 'Valencia Late' sweet oranges grown on three rootstocks in eastern Mediterranean

Ercan Yildiz^{1*}, Turan Hakan Demirkese¹, and Mustafa Kaplankiran¹

The plant growth, yield, yield efficiency, and fruit quality of 'Rhode Red Valencia' and 'Valencia Late' sweet oranges (*Citrus sinensis* (L.) Osbeck) on three rootstocks were evaluated under Eastern Mediterranean climatic conditions of Dörtol-Hatay, which is one of the oldest and largest citrus and mandarin (*Citrus reticulata* Blanco) production regions in Turkey. The fruit yield was affected by rootstock in both scion cultivars from 2007 through 2010. Trees on 'Troyer' citrange had lower yield than those budded on the other rootstocks. The yield of 'Rhode Red Valencia' and 'Valencia Late' orange trees on 'Carrizo' citrange were about 9% and 19% more than those on 'Troyer' citrange, respectively. But trees on 'Troyer' citrange had significant higher yield efficiency than trees on sour orange and 'Carrizo' citrange, because of canopy volume of 'Troyer' citrange was lower than the other rootstocks. The heaviest fruits of 'Valencia Late' orange trees were harvested from 'Carrizo' citrange (214.69 g), while 'Rhode Red Valencia' orange trees did not show differences regarding the rootstock. The rootstocks had no significant effects on juice content of 'Rhode Red Valencia' and 'Valencia Late' oranges. The effects of the rootstocks in both scion cultivars on juice content, total acids (TA), Brix:TA ratio, and number of seeds per fruit were found to be insignificant.

Key words: Orange, rootstock, fruit quality, plant characteristics, yield.

INTRODUCTION

Turkey's citrus production has reached to approximately 3 600 000 t in 2010; with an increase of 44.2% in the past decade. The statistics of 2010 indicated that among the Turkey's total citrus production, orange is the first (1 710 500 t) species (FAOSTAT, 2010).

The 'Valencia Late' orange (*Citrus sinensis* (L.) Osbeck) is one of the sweet oranges used for juice extraction. It is a late-season fruit, and therefore a popular variety when the navel oranges are out of season. The popularity of 'Valencia Late' juice has been stimulated by the market success of the not-from-concentrate (NFC) pasteurized product. Juice color of the 'Valencia' clone means less juice needed for blending to attain the desired color. 'Rhode Red Valencia' is a clone that has been selected for its high juice color score (Rouse, 2000).

Due to the changes in customer preferences and demands, many new citrus cultivars have been introduced in Turkey. It is not expected that the introduced cultivars will perform similarly in various regions of Turkey where different ecological conditions exist. The different ecological condition effects on citrus productions are apparent. Hence, it is valuable to know the favorable

ecological conditions for the cultivars chosen and their interactions under these parameters by ecological conditions of the growing sites. Further, factors like cultivar characteristics, rootstocks employed, growing conditions along with cultural managements, type of flowers, and the fruit drops can affect citrus cultivars yield and quality performance (Demirkese et al., 2003). Among the cultivars that are mostly used in new orchards of Turkey, 'Okitsu' and 'Clausellina' have been planting in the orchards established since the beginning and end of 1990s, respectively (Cinar, 2004). Kaplankiran et al. (2005) suggested avoiding the over-production during the harvest period when the prices are low; growers should focus on the 'Okitsu' and 'Clausellina' mandarins and 'Midnight Valencia' and 'Rhode Red Valencia' oranges.

Rootstocks have had a substantial role in the development of the citrus industry in the world. Rootstock utilization has gain value for solving both limiting and restricting factors of citrus production (soil, climate and pests, etc.) and conditioning the market demands on productivity, short juvenility period and high-fruit quality (Tuzcu et al., 1992). The rootstock may influence several aspects of citrus growth and development, including yield, fruit quality, and tolerance to stress caused by biotic and abiotic factors (Filho et al., 2007). The main rootstock in Turkey for citrus production is 'Sour Orange', while trifoliate orange, 'Troyer' and 'Carrizo' citrange are also used. The use of 'Carrizo' has been increased lately, especially in Eastern Mediterranean Region (Kaplankiran et al., 2001).

¹University of Mustafa Kemal, Faculty of Agriculture, 31034 Antakya, Hatay, Turkey. *Corresponding author (ercanyildiz54@gmail.com).

Received: 14 November 2012.

Accepted: 4 March 2013.

doi:10.4067/S0718-58392013000200009.

This study was conducted to determine and compare the yield and fruit quality performances of three rootstocks for ‘Rhode Red Valencia’ and ‘Valencia Late’ in Eastern Mediterranean region of Turkey.

MATERIALS AND METHODS

The used rootstocks were ‘Sour orange’ (*Citrus aurantium* L. ‘Yerli’) and ‘Carrizo’ and ‘Troyer’ citranges (*Poncirus trifoliata* Raf. × *Citrus sinensis* (L.) Osb. ‘Carrizo’ and ‘Troyer’). Trees were planted in 1998 at the Research Station of Mustafa Kemal University, Dörtöyl (36°09’ E; 36°51’ N; 9 m a.s.l.), Turkey. The experimental design was a randomized complete blocks design with five replicates and a single-tree per plot. Trees were 7 m × 7 m apart under drip irrigation system and received the same cultural practices.

The soil texture of the plot located in Dörtöyl (Hatay) was sandy-silt texture. The soil was course textured (17.6% coarse sand, 37.6% fine sand, 23.8% silt, 22% clay) and slightly alkaline to alkaline in the soil profile (pH 7.80, 7.98 and 8.25 for 0-30, 30-60 and 60-90 cm depth, respectively, in 1:2.5 soil:water suspension), rich in carbonate content (61-63 g kg⁻¹ for 0-60 cm and 113.5 g kg⁻¹ for 60-90 cm depth).

In January of each season, tree height, canopy diameter in the two tree directions (to obtain the average diameter), canopy circumference, trunk girth at 10 cm above and below the budding union were measured, and scion/stock girth ratio was calculated. The canopy volume (CV) was calculated from canopy height and spread, considering canopy as a prolate spheroid and applying the formula: CV = 4/3 πab² where a is the major axis length/2, and b is the minor axis length/2 (Westwood, 1993).

Fruits per tree were collected and separately weighed so that the yield per tree (kg) could be calculated. The ratio of yield to CV (yield efficiency, kg m⁻³) and the cumulative yield/tree were also calculated. Alternate bearing index (ABI) was calculated between the years 2007 and 2010, using the following expression (Monselise and Goldschmidt, 1982):

ABI = 1 / (n - 1) × { (Ia2 - a1I) / (a2 + a1) + (Ia3 - a2I) / (a3 + a2) + ... + (Ia_n - a_{n-1}I) / (a_n + a_{n+1}) }

where n is number of years, and a1, a2, ..., a_{n-1}, a_n are yields of the corresponding years.

Samples of 20 fruits per replicate were randomly collected at harvest date for determining physical and chemical characteristics. Individual fruit weight, fruit diameter, rind thickness, number of seeds per fruit, juice content, total soluble solids or °Brix, titratable acidity (TA) and then °Brix/TA ratio were determined.

Data recorded in all seasons were subjected to ANOVA using SAS program (SAS Institute, 1999) and means were compared with Tukey test at 5% level of significance.

RESULTS AND DISCUSSION

Plant growth

At 12 yr-old, trunk cross-sectional area (TCSA) and canopy projection area of both ‘Rhode Red Valencia’ and ‘Valencia Late’ orange trees on ‘Troyer’ citrange were significantly lower than those on ‘Carrizo’ citrange and sour orange (Table 1). Similar results were reported by Georgiou and Gregoriou (1999) on ‘Shamouti’ orange, who stated that ‘Troyer’ citrange induced the lowest values of TCSA as compared with ‘Carrizo’ citrange and sour orange, while Georgiou (2002; 2004) found that TCSA of ‘Clementine’ mandarin and ‘Valencia’ orange trees budded on ‘Carrizo’ and ‘Troyer’ citranges was similar, and lower than those on sour orange.

The largest canopy height of ‘Valencia Late’ orange trees were on ‘Carrizo’, although canopy heights of ‘Rhode Red Valencia’ orange trees were not significantly different according to rootstocks. Canopy diameters of both scion cultivars trees on ‘Carrizo’ were significantly larger than other rootstocks (Table 1). The results agreed with ‘Carrizo’ rootstock promoted higher canopy height and diameter than other rootstocks on ‘Orlando’ tangelo (Fallahi et al., 1991).

The canopy volume (CV) of ‘Rhode Red Valencia’ and ‘Valencia Late’ on ‘Carrizo’ were approximately 34% and 12% higher, respectively, than those on ‘Troyer’ (Table 1). Our CV findings are in harmony with those of Hutchison and Hearn (1977) on ‘Robinson’ and ‘Page’ mandarin trees; and those of Euler et al. (2008) on ‘Valencia’ orange

Table 1. Effect of rootstock on some plant characteristics of different “Valencia” clone trees in 2010.

Cultivar	Rootstock	Trunk cross-sectional area	Canopy projection area	Canopy height	Canopy diameter	Canopy volume	Scion:stock ratio ¹
		cm ²	m ²	m		m ³	
Rhode Red Valencia	Sour orange	234.37a	11.58ab	3.34	3.84ab	24.29	0.92
	Carrizo	243.27a	12.56a	3.40	4.00a	26.55	0.89
	Troyer	197.53b	10.76b	3.40	3.70b	23.65	0.89
HSD		18.53	1.27	NS	0.18	NS	NS
Valencia Late	Sour orange	242.55a	12.27a	3.32ab	3.93ab	25.88a	0.90
	Carrizo	242.05a	12.69a	3.43a	4.01a	27.10a	0.85
	Troyer	172.61b	10.62b	3.04b	3.67b	20.25b	0.89
HSD		39.14	1.45	0.28	0.26	4.10	NS

Means in the same column followed by different letters are significantly different according to Tukey’s New Multiple Range test (p ≤ 0.05). NS: Non significant; HSD: Honestly significant difference.

¹Ratio of scion trunk circumference to rootstock trunk circumference.

trees budded on different rootstocks, who found that the lowest CV was with ‘Troyer’. Further, Gonzalez-Velez et al. (2002) mentioned that CV and canopy height of ‘Orlando’ tangelo trees was not affected by the rootstocks. These characteristics of rootstocks must be considered to determine in row and spacing of the row in establishing the orchard.

Rootstock influence on scion:stock ratio was statistically similar, although the most favorable rootstocks for ‘Rhode Red Valencia’ and ‘Valencia Late’ were sour orange (Table 1). Similar results were also obtained on ‘Valencia’ (Hassan et al., 2000) and ‘Shamouti’ oranges (Georgiou and Gregoriou, 1999) and ‘Nova’ (Georgiou, 2000), ‘Clementine’ (Georgiou, 2002) and ‘Marisol’ mandarins (Bassal, 2009).

Yield

The fruit yield was affected by the rootstock in both scion cultivars, from 2007 through 2010, except for ‘Valencia Late’ in 2009. Trees on ‘Troyer’ had lower yield than those budded on the other rootstocks. The yield of ‘Rhode Red Valencia’ and ‘Valencia Late’ orange trees on ‘Carrizo’ were about 9% and 19% more than those on ‘Troyer’, respectively. Yield of the trees on sour orange were found between ‘Carrizo’ and ‘Troyer’ rootstocks (Table 2). Similar results were also obtained by Georgiou (2002), who found that mandarin trees on ‘Carrizo’ were more productive than those on ‘Troyer’. On the contrary, ‘Olinda Valencia’ (Al-Jaleel and Zekri, 2002) and ‘Parent Washington Navel’ (Al-Jaleel and Zekri, 2003) oranges were reported to yield similarly on sour orange and ‘Carrizo’ rootstocks. ‘Carrizo’ rootstock promoted higher yield compared to other rootstocks (Rouse, 2003; Tuzcu et al., 2004; Kaplankiran et al., 2005). Further, Filho et al. (2007) mentioned that fruit yield of ‘Fallglo’ and ‘Sunburst’ mandarin trees was not affected by the rootstock. All these results indicated the inconsistency in yield differences as affected by rootstocks, which could be attributed to differences in scion cultivars, tree age, climatic conditions, and soil characteristics, etc.

Having lower CV, trees on ‘Troyer’ had significantly higher yield efficiency than trees on sour orange and

‘Carrizo’ (Table 2). The lower yield efficiency on sour orange and ‘Carrizo’ is probably caused by the bigger canopy size induced by these rootstocks. The higher yield efficiencies were also reported for trees reduced in size by rootstocks (Roose et al., 1989; Auler et al., 2008).

‘Rhode Red Valencia’ orange trees on ‘Troyer’ showed significant ABI, however, no significant differences were found among these rootstocks for ‘Valencia Late’ (Table 2), which is supported by previous studies (Georgiou, 2000; 2002; Filho et al., 2007). The ABI may be reduced with the control of the crop load during the ‘on’ years by cultural practices, such as fruit thinning, branch girdling, exogenous application of plant growth regulators, and early harvest (Sposito et al., 1998).

Fruit quality

The juice content, total acids (TA), °Brix:TA ratio and number of seeds per fruit were not affected by the rootstock, in both scion cultivars. The fruit weight was higher in fruits of ‘Valencia Late’ trees budded on ‘Carrizo’ (214.69 g), while ‘Rhode Red Valencia’ trees (between 180.22 and 186.98 g) did not show any differences regarding the rootstock for this variable (Table 3). The results for ‘Rhode Red Valencia’ are in agreement with those of Myhob et al. (1996) on ‘Valencia’ orange; Gonzalez-Velez et al. (2002) on ‘Orlando’ tangelo, and Filho et al. (2007) on ‘Fallglo’ and ‘Sunburst’ mandarins, who reported that fruit weight was not affected by the rootstock. Auler et al. (2008) on ‘Valencia’ orange trees budded on six rootstocks declared that the trees on ‘Troyer’ produced smaller fruit than those on other rootstocks. Al-Jaleel and Zekri (2002) on ‘Olinda Valencia’ orange trees budded on nine rootstocks; and Auler et al. (2008) on ‘Valencia’ orange trees budded on six rootstocks reported that the fruit weight was determined in the range of 188.4-239.8 and 164.2-183.7 g, respectively.

The higher values of fruit diameter were verified in fruits of ‘Rhode Red Valencia’ orange trees on ‘Carrizo’ and ‘Troyer’, however ‘Valencia Late’ on ‘Carrizo’ had the least fruit diameter. The highest rind thickness was also obtained in fruits of ‘Rhode Red Valencia’ on ‘Troyer’ and those of ‘Valencia Late’ on ‘Carrizo’ and ‘Troyer’ (Table 3).

Table 2. Effect of rootstock on yield according years, cumulative yield, yield efficiency (yield in 2010 per tree volume) and ABI of different “Valencia” clone trees.

Cultivar	Rootstock	Yield (kg per tree)				Cumulative 2007-2010	Yield efficiency	ABI
		2007	2008	2009	2010			
Rhode Red Valencia	Sour orange	84.2a	81.0a	126.0ab	145.2b	436.4a	6.06b	0.11b
	Carrizo	84.4a	81.0a	109.4b	164.4a	439.2a	6.21ab	0.18ab
	Troyer	59.8b	56.3b	134.0a	152.6ab	402.7b	6.70a	0.26a
HSD		15.06	15.53	20.19	14.04	26.54	0.58	0.08
Valencia Late	Sour orange	42.5a	70.8a	100.4	165.2b	378.9a	7.15b	0.27
	Carrizo	45.3a	68.5a	107.2	182.0a	403.1a	6.80b	0.25
	Troyer	26.8b	50.5b	97.8.	162.4b	337.5b	8.68a	0.28
HSD		5.99	11.95	NS	10.69	27.87	0.96	NS

Means having the different letters in each column are significantly according to Tukey test (p ≤ 0.05).
NS: Non significant; ABI: Alternate bearing index; HSD: Honestly significant difference.

Table 3. Effect of rootstock on fruit quality of different "Valencia" clone (Average 2007-2010).

Cultivar	Rootstock	Fruit weight g	Fruit diameter mm	Rind thickness	Juice content	°Brix %	Total acids (TA)	°Brix: TA ratio	Number of seeds per fruit
Rhode Red Valencia	Sour orange	180.22	69.57b	4.01b	50.60	9.09a	1.29	7.12	3.38
	Carrizo	186.98	72.59a	4.16b	50.09	9.00a	1.23	7.20	3.51
	Troyer	185.36	72.05a	4.53a	49.17	8.82b	1.25	7.14	3.01
HSD		NS	2.42	0.29	NS	0.17	NS	NS	NS
Valencia Late	Sour orange	195.84b	75.25b	3.98b	47.88	9.23	1.37	6.76	4.78
	Carrizo	214.69a	80.15a	4.85a	49.23	9.22	1.36	6.77	4.77
	Troyer	187.68b	77.11b	4.69a	48.24	9.14	1.34	6.80	4.74
HSD		9.40	2.54	0.50	NS	NS	NS	NS	NS

NS: Non significant; HSD: Honestly significant difference.

Means having the different letters in each column are significantly according to Tukey test ($p \leq 0.05$).

The rootstocks had no significant effects on juice content of 'Rhode Red Valencia' and 'Valencia Late', which is ranged between 49.17-50.60% and 47.88-49.23%, respectively. Gregoriou and Economides (1993) on 'Ortanique' tanger; Tuzcu et al. (2004) on 'Washington Navel' orange; and Filho et al. (2007) on 'Fallglo' and 'Sunburst' mandarins mentioned that fruit juice content was not affected by the previous studies on rootstock. Al-Jaleel and Zekri (2002) on 'Olinda Valencia' orange in Saudi Arabia and Auler et al. (2009) on 'Valencia' orange in Brazil conditions reported that fruits on nine rootstocks had juice content ranged between 51.1-53.7% and 48.8-51.3%, respectively.

Although the °Brix was not significantly different by the rootstocks on 'Valencia Late', it was lower in fruits of 'Rhode Red Valencia' on 'Troyer' (Table 3). The results obtained by 'Valencia Late' are in agreement with those of Hearn (1987) on 'Fallglo' mandarin and Hearn (1989) on 'Ambersweet' orange, who found that the fruits on 'Carrizo' had similar °Brix those on sour orange. °Brix findings provided from this study were lower than the results of Al-Jaleel and Zekri (2002) on 'Olinda Valencia' and Auler et al. (2009) on 'Valencia' oranges reported that fruits on all rootstocks had Brix between 11.8% and 13.7%, and between 10.3% and 11.4%, respectively.

The total acids of both 'Rhode Red Valencia' and 'Valencia Late' orange trees were not affected by the rootstocks (Table 3). Myhob et al. (1996) on 'Valencia'; Tuzcu et al. (2004) on 'Washington Navel' orange; Demirkese et al. (2005) on 'Rhode Red Valencia' orange; Kaplankiran et al. (2005) on 'Okitsu' Satsuma mandarin; and García-Sánchez et al. (2006) on 'Clemenules' mandarin found that the effects of the rootstocks on fruit juice acidity were insignificant.

CONCLUSIONS

Both scion cultivars displayed high yield efficiency and weak growth on 'Troyer' citrange, which may be better suited to high density plantings. On the other hand, 'Carrizo' citrange and sour orange seems to be more suitable rootstock in terms of yield per tree for

the cultivars used in this study. Fruit quality of Rhode Red Valencia was similar for each rootstock used in this experiment, but it was higher on 'Carrizo' citrange in 'Valencia Late' orange. At present in Turkey, sour orange is the most commonly used rootstock for the commercial cultivation of 'Valencia Late' orange but it is highly susceptible to *Citrus tristeza virus*. The results showed that 'Carrizo' and 'Troyer' citranges, which is supported by previous research, were very promising as alternative rootstocks. However, further studies in different locations with different conditions are needed, and the current study should be prolonged to see the effects age of the tree.

LITERATURE CITED

- Al-Jaleel, A., and M. Zekri. 2002. Yield and fruit quality of 'Olinda Valencia' trees grown on nine rootstocks in Saudi Arabia. *Proceedings of the Florida State Horticultural Society* 115:17-22.
- Al-Jaleel, A., and M. Zekri. 2003. Effects of rootstocks on yield and fruit quality of 'Parent Washington Navel' trees. *Proceedings of the Florida State Horticultural Society* 116:270-275.
- Auler, P.A.M., A.C.G. Fiori-Tutida, and M.B. dos S. Scholz. 2009. Industrial quality and maturation of fruits of 'Valencia' sweet orange trees on six rootstocks. *Revista Brasileira de Fruticultura* 31(4):1158-1167.
- Auler, P.A.M., A.C.G. Fiori-Tutida, and Z.H. Tazima. 2008. Behavior of 'Valencia' orange tree on six rootstocks in the northwest of Paraná state. *Revista Brasileira de Fruticultura* 30(1):229-234.
- Bassal, M.A. 2009. Growth, yield and fruit quality of 'Marisol' clementine grown on four rootstocks in Egypt. *Scientia Horticulturae* 119(2):132-137.
- Cinar, A. 2004. The structure of citrus cultivation in Turkey and some developments in recent years (in Turkish with an English summary). *Cine Tarım* 7(57):14-17 (Tr).
- Demirkese, T.H., S. Eti, and M. Kaplankiran. 2003. The effects of GA₃ and BA treatments on fruit set and quality on Nova mandarin (in Turkish with an English summary). p. 181-184. IV National Symposium on Horticulture (Tr), Antalya, Turkey. 8-12 September. Akdeniz University, Antalya, Turkey.
- Demirkese, T.H., M. Kaplankiran, and E. Yildiz. 2005. The effects of some citrus rootstocks on fruit yield and quality for Rhode Red Valencia orange during the periode of the juvenility in Dörtöl (Hatay- Turkey) conditions. p. 27-28. *Book of Abstracts of Seventh International Congress of Citrus Nurserymen*, Cairo, Egypt. 17-21 September 2005. Cairo University, Cairo, Egypt.
- Fallahi, E., Z. Mousavi, and D.R. Rodney. 1991. Performance of 'Orlando' tangelo trees on ten rootstocks in Arizona. *Journal of the American Society for Horticultural Science* 116:2-5.

- Filho, F.A.A.M., E. Espinoza-Núñez, E.S. Stuchi, and E.M.M. Ortega. 2007. Plant growth, yield, and fruit quality of 'Fallglo' and 'Sunburst' mandarins on four rootstocks. *Scientia Horticulturae* 114:45-49.
- FAOSTAT. 2010. FAO statistical databases. Agriculture. Available at <http://faostat.fao.org/site/339/default.aspx> (accessed 4 June 2012).
- García-Sánchez, F., J.G. Perez-Perez, P. Botia, and V. Martínez. 2006. The response of young mandarin trees grown under saline conditions depends on the rootstock. *European Journal of Agronomy* 24:129-139.
- Georgiou, A. 2002. Performance of 'Nova' mandarin on eleven rootstocks in Cyprus. *Scientia Horticulturae* 84:115-126.
- Georgiou, A. 2002. Evaluation of rootstocks for 'Clementine' mandarin in Cyprus. *Scientia Horticulturae* 93:29-38.
- Georgiou, A. 2004. Evaluation of rootstocks for 'Valencia' orange. *Agricultura Mediterranea* 134:193-200.
- Georgiou, A., and C. Gregoriou. 1999. Growth, yield and fruit quality of 'Shamouti' orange on fourteen rootstocks in Cyprus. *Scientia Horticulturae* 80:113-121.
- Gregoriou, C., and C.V. Economides. 1993. Tree growth, yield, and fruit quality of Ortanique tangor on eleven rootstocks in Cyprus. *Journal of the American Society for Horticultural Science* 118:335-338.
- Gonzalez-Velez, A., Roman-Perez, F., and C. Flores. 2002. Tangelo 'Orlando' grafted on five rootstocks: production, fruit quality and growth. *Journal of Agriculture of the University of Puerto Rico* 86(3/4):131-137.
- Hassan, A.S., S.I. Gaafer, M.H. Saad-Allah, and A.M. Ibrahim. 2000. Effect of some citrus rootstocks on growth of young Baladi mandarin and Valencia orange trees in newly reclaimed soil. II. Growth flushes and leaf chemical constituents. *Zagazig Journal of Agricultural Research* 27(4):965-989.
- Hearn, C.J. 1987. The Fallglo citrus hybrid in Florida. *Proceedings of the Florida State Horticultural Society* 100:119-121.
- Hearn, C.J. 1989. Yield and fruit quality of 'Ambersweet' orange hybrid on different rootstocks. *Proceedings of the Florida State Horticultural Society* 102:75-78.
- Hutchison, D.J., and C.J. Hearn. 1977. The performance of Nova and Orlando tangelos on 10 rootstocks. *Proceedings of the Florida State Horticultural Society* 90:47-49.
- Kaplankiran, M., T.H. Demirkaser, C. Toplu, and M. Uysal. 2001. The structure of citrus production, the status of rootstocks and nursery tree production in Turkey. p. 190-194. *Proceeding of 6th International Congress of Citrus Nurserymen*, Ribeirao Preto, Brazil. 9-13 July 2001. Sao Paulo University, Ribeirao Preto, Brazil.
- Kaplankiran, M., T.H. Demirkaser, and E. Yildiz. 2005. The effects of some citrus rootstocks on fruit yield and quality for Okitsu Satsuma during the juvenility period in Doryol (Hatay, Turkey) conditions. p. 27. *Book of Abstracts of Seventh International Congress of Citrus Nurserymen*, Cairo, Egypt. 17-21 September 2005. Cairo University, Cairo, Egypt.
- Monselise, S.P., and E.E. Goldschmidt. 1982. Alternate bearing in fruit trees. p. 128-166. In J. Janic (ed.) *Horticultural reviews*. AVI Publishing Company, Westport, Connecticut, USA.
- Myhob, M.A., M.H. Saad-Allah, T.A. Ibrahiem, and S.E. Salem. 1996. Evaluation of 'Valencia' orange trees grafted on sour orange, Succari orange and Troyer citrange. *Bulletin of Faculty of Agriculture, Cairo University* 47(3):449-456.
- Roose, M.L., P.A. Cole, D. Atkin, and R.S. Kupper. 1989. Yield and tree size of four citrus cultivars on 21 rootstocks in California. *Journal of the American Society for Horticultural Science* 114:678-684.
- Rouse, R.E. 2000. Citrus fruit quality and yield of six Valencia clones on 16 rootstocks in the immokalee foundation grove. *Proceedings of the Florida State Horticultural Society* 113:112-114.
- Rouse, R.E. 2003. Production, fruit quality, and color development of Rhode Red Valencia oranges on different rootstocks in subtropical Florida. *Proceedings of the Interamerican Society for Tropical Horticulture* 47:248-251.
- SAS Institute. 1999. *SAS/STAT User's guide*. SAS Institute, Cary, North Carolina, USA.
- Sposito, M.B., P.R.C. Castro, and M. Agusti. 1998. Alternância de produção em citros. *Laranja* 19:293-304 (Fr).
- Tuzcu, O., M. Kaplankiran, S. Duzenoglu, T. Bahceci, and T. Yesiloglu. 1992. Effect of some citrus rootstocks on the yield and quality of the 'Washington Navel' orange variety in Adana (Turkey) conditions. p. 270-274. *7th International Citrus Congress*, Acireale, Italy. 8-13 March 1992. Agricultural Research Council (CRA), Acireale, Italy.
- Tuzcu, O., B. Yildirim, and T. Yesiloglu. 2004. Effect of different rootstocks and sectors on fruit yield and its distribution depending to the tree canopy. p. 515-519. *10th International Society of Citriculture Congress*, Agadir, Morocco. 15-20 February 2004. University of Agadir (Ibn Zohr), Agadir, Morocco.
- Westwood, M.N. 1993. *Temperate-zone pomology: Physiology and culture*. 3rd ed. 535 p. Timber Press, Portland, Oregon, USA.

Copyright of Chilean Journal of Agricultural Research is the property of Instituto de Investigaciones Agropecuarias (INIA) and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.