

***In Situ* Localization of Germin Gene Expression During Auxin Induced Callus Formation**

Mahmut ÇALIŞKAN

Mustafa Kemal University, Department of Biology
31040 Hatay - TURKEY

Received: 21.01.2000

Abstract: Wheat embryogenic callus was induced by treating immature wheat embryos with 2,4-D. Digoxigenin-labeled germin sense and anti-sense RNA probes were prepared by *in vitro* transcription. By using these probes it was revealed that there was a striking induction of germin gene expression prior to visible callus formation. Although no germin gene expression was localized on the sections of immature embryos which were not incubated in callus induction medium, following 4 hours incubation in callus induction medium, germin mRNA was localized on the sections of immature embryos. Germin gene expression more conspicuous and widespread on the sections which were incubated for 6 hours in callus induction medium. It was suggested that germin genes induced during callus formation were related to the remodeling of the cell wall to accomplish the transition from precocious germination to callus formation.

Key Words: Callus, Germin, Auxin, Embryogenesis

Oksin'in Oluşturduğu Kallus Dokusunda Germin Gen Ekspresyonunun Belirlenmesi

Özet: Olgunlaşmamış buğday embriyoları 2, 4-D ile muamele ettirilerek embriyonik kallus oluşturulmuştur. Germin'in digoxigenin ile işaretlenmiş anlamlı ve anlamsız RNA problemleri laboratuvar koşullarında hazırlanmıştır. Bu problemler kullanılarak, oksinle muameleye tabi tutulan embriyolarda, belirgin kallus oluşumundan önce önemli oranda germin gen ekspresyonunun olduğu anlaşılmıştır. Oksinsiz ortamda bulunan embriyolarda germin gen ekspresyonu belirlenmemesine rağmen, oksin'li ortamda 4 saatlik bir inkübasyon sonucunda embriyolarda, germin gen ekspresyonunun olduğu belirlenmiştir. Germin gen ekspresyonunun, oksinli ortamda 6 saatlik bir inkübasyon sonunda, çok daha yaygın ve belirgin olduğu gözlemlenmiştir. Germin gen ekspresyonlarının, olgunlaşmamış buğday embriyolarının erken çimlenme aşamasından kallus oluşumu aşamasına geçişi gerçekleştirmek için gerekli olan, hücre çeperinin yeniden şekillenmesi ile ilgili olabileceği önerilmiştir.

Anahtar Sözcükler: Kallus, Germin, Oksin, Embriyogenez

Introduction

Wheat embryo development is initiated by fertilization of the ovum, following which the formation of a viable embryo passes through several distinct stages. Following a period of cellular proliferation and differentiation, during which the primordial organs are formed, the embryo achieves a state of functional maturity such that its dissection from the developing grain permits the onset of precocious germination (1,2). If left within the grain, the embryo remains in a dormant state; a stage during which the accumulation of a discrete set of maturation-associated gene products occurs (3,4). The embryogenic phase of development is brought to a close by the programmed desiccation of the grain, causing a cessation of metabolic activity in the embryo. Desiccation causes a change to occur in the developmental potential of the embryo so that upon rehydration, instead of resuming an anabolic embryogenic pathway of development, the embryonic metabolism switches into a catabolic germinative mode utilizing embryonic and endosperm reserves to fuel the explosive growth of root and shoot meristems.

The programming of embryonic development is, in part, regulated by plant growth substances such as auxin, abscisic acid (ABA), gibberellic acid (GA), and ethylene (5-9). This also is associated with a change in the genetic program. For instance, successful germination requires the transcription of new genes (10). Conspicuous amongst the population of germination associated gene products is the remarkable homopentameric, extra cellular glycoprotein called "germin" (11,12). Germin genes have been shown to be associated with various aspect of plant development such as salt stress (13,14), defense system (15-18), embryonic development (19), photoperiodic oscillations (20, 21) and hormonal stimuli (22). Germin was suggested to be a member of a "superfamily" which comprises various growth related genes (23-25). Furthermore, wheat germin gene expression was shown to be regulated by biotic (viral infection) and abiotic stress (heavy metal ions) applications (26).

In 1993, it was determined that cereal germin proteins had strong oxalate oxidase activity (27,28). Oxalate oxidase (EC 1.2.3.4) is an oxidoreductase which degrades oxalic acid to produce one mole of H_2O_2 and two moles of CO_2 . The principal product of the degradation of oxalate is hydrogen peroxide, which is a highly reactive compound known to be involved in several metabolic processes in higher plants. It was reported that hydrogen peroxide might act as a signaling molecule at low concentrations (29) or a component of cell wall modifications at high concentrations (30). The involvement of oxalate, germin genes and hydrogen peroxide in plant development was reviewed by Caliskan (31,32). The role of germin genes and their enzymatic activity during wheat embryo germination have been studied (33-35). Our previous results revealed that germin expression was associated with the tissues have caused whose growth was restricted during germination. It was reported that germin might have caused the growth restriction by inducing cross-linking of cell wall polymers prompted by H_2O_2 accumulation, which is a product of germin action (33).

Culture of immature wheat embryos, isolated from developing grains, in the presence of 2,4-D caused scutellar tissue to undergo unchecked proliferation, as callus (36). Callus has the capacity to undergo subsequent regeneration via somatic embryogenesis, a process of some significance for the potential genetic manipulation of this important cereal (36,37). Since sequence analysis of a wheat germin gene indicated the presence of potentially auxin responsive *cis*-acting elements in the promoter region (38), the expression pattern of germin genes in callus formation has been investigated in the current study in order to characterize the processes inherent in the capacity of embryonic wheat tissues for dedifferentiation and redifferentiation.

Materials and Methods

Plant material

Grains of spring wheat (*Triticum aestivum* L. var. Tonic) obtained from Kenneth Wilson Grain, Leeds, UK, were grown in a controlled-environment chamber with a 16h light (25°C) - 8 h, dark (18°C) photoperiod at 60% relative humidity. Immature grains were harvested at 12 days after fertilization (12 daf). For callus induction, grains were surface sterilized with sodium hypochlorite solution (1% free Cl₂) for 10 minutes and washed with sterile water prior to dissection of the embryos. Embryos were washed in sterile water to remove adhering liquid endosperm and transferred to 0.6% agar medium containing Murashige-Skoog medium (MS) supplemented with 2% (w/v) sucrose with or without 2mg l⁻¹ 2,4-dichlorophenoxyacetic acid (2,4-D; Sigma, UK) (36). Cultures were incubated at 25°C in darkness until use. Sections were prepared from cultures following 0-hour (immediately), 4-hour and 6-hour incubation periods in callus induction medium.

In situ hybridization

"Germin" mRNA was detected in sections by hybridization, *in situ*, with transcripts of a "germin" cDNA, kindly provided by Prof. Dr. B. G. Lane (University of Toronto). This cDNA was subcloned in the plasmid vector pBluescript (Stratagene, La Jolla, CA) for the production of digoxigenin-labeled riboprobes (DIG system: Boehringer Mannheim, FRG). Labeled anti-sense and sense probes were produced with T3 and T7 RNA polymerases (Figure 1). Probes were subjected to mild alkaline hydrolysis by incubation with 40mM NaHCO₃, pH 10.2 at 60°C to produce fragments of *ca.* 250 bp (39). Subsequent experiments showed that labeled anti-sense riboprobes specifically hybridized with germin mRNA in tissues, while labeled sense riboprobes gave no hybridization product (data not shown). Therefore, labeled sense riboprobes were used in control experiments in which no signal was expected.

A modified procedure of Jackson (40) was employed for localization of mRNA *in situ*. Samples were prepared for sectioning by fixation in 4% (w/v) paraformaldehyde in phosphate buffered saline (PBS) at 4°C, dehydrated through ethanol series on ice, and embedded in




Figure 1. Production of RNA probes. The germin cDNA was cloned in the Eco RI site of pBluescript II SK⁺. The plasmid DNA was linearized with Bam HI for production of T7 transcripts (sense) and Eco RV for production of T3 transcripts (anti-sense).

paraffin wax using Histolene™ (CellPath, Hemel Hempsted, UK) as the infiltrating solvent. Sections (10µm) were mounted on poly-L-lysine (Sigma, UK) coated slides at 42°C overnight, deparaffinized and rehydrated through ethanol series for hybridization with strand-specific probes. The sections were incubated with pronase (0.125mg ml⁻¹ in 50mM Tris-Cl - 5mM EDTA, pH 7.5) and post-fixed with 4% (w/v) paraformaldehyde in PBS. Sections were acetylated with acetic anhydride (0.5% (v/v) in triethanolamine-HCl, pH 8) prior to prehybridization at room temperature with 0.3M NaCl - 10mM sodium phosphate buffer - 10mM Tris-Cl - 5mM EDTA - 50% (v/v) formamide - 10% (w/v) dextran sulphate - 0.1% (w/v) tRNA - 1x Denhardt's solution, pH 6.8 for 30 minutes. Sections were then incubated with buffer containing the DIG-labeled probe at a final concentration of 3 µg ml⁻¹ overnight at 50°C. Sections were washed in 2xSSC-50% (v/v) formamide at 50°C, and then incubated with RNase A (20 µg ml⁻¹) in 0.5M NaCl, 10mM Tris-Cl, pH 7.7, 1mM Na₂EDTA at 37°C for 30 min. After a final wash in PBS, the hybridized probe was detected using anti-digoxigenin antiserum at a 1:3000 dilution, as described in the manufacturer's instructions. Sections were examined by light microscopy and photographed with Kodak Ektachrome Elite II color film.

Results and Discussion

Immature wheat embryos excised from developing grains and cultured in the presence of auxin do not continue their normal pathway of development. Instead, cells of the immature embryos become dedifferentiated and proliferate to form unchecked cell growth that is called callus. Although germin synthesis was not caused by either immature embryos or mature embryos before their germination (34,41), an abundant accumulation of germin gene expression was detected during callus formation from auxin incubated immature wheat embryos (41). The localization pattern of germin gene expression in a spatial and temporal manner during callus formation was the main concern of the current study.

Immature wheat embryos have a conspicuous appearance of primary and secondary roots, scutellum, coleorhiza and coleoptile structures at 12 days post-anthesis (Figs. 2A, 2B). The first two sections were taken from the immature embryos, which were not incubated in callus induction medium, and they were called 0-hour. When these sections were reacted with labeled anti-sense germin riboprobes, no hybridization product was observed (Figure 2A). The sections reacted with labeled sense germin riboprobe also gave no signals as expected (Figure 2B). This was completely in accordance with previous studies, which reported that there was no germin gene expression in immature wheat embryos (41). Following 4-hour incubation in callus induction medium, however, the immature embryo sections revealed some germin gene




Figure 2. Immature wheat embryo sections treated with germin DIG-labeled (A) anti-sense RNA probes and (B) sense RNA probes (0-hour, no incubation in callus induction medium). S: Scutellum, C: Coleorhiza, PR: Primary root.

expression signals, particularly on the outer cells of the scutellum (Figure 3A). In 4 hours, callus induction medium caused expression of germin genes in the scutellum, which is an embryonic tissue responsible for protecting and supporting the embryo development. The sections taken from the same embryo were reacted with labeled sense riboprobes for the control experiments and no signal was observed (Figure 3B). Therefore, the signals associated with the scutellum were really a product of hybridization between labeled anti-sense germin riboprobes and germin mRNA in cells. Germin gene expression was more conspicuous when immature wheat embryos were incubated for 6 hours in callus induction medium. At that stage of development, germin gene expression was found to be associated with outer scutellum cells, particularly the tips of the scutellum, and the tips of coleorhiza cells (Figure 4A). Although there was some background staining on the primary root in the control section, which was reacted with labeled sense riboprobes (Figure 4B), it is obvious from the figure that primary root cells gave some signals of germin gene expression (Figure 4A). It was observed that there was no visible callus




Figure 3. Immature wheat embryo sections treated with germin DIG-labeled (A) anti-sense RNA probes and (B) sense RNA probes (after 4-hour incubation in callus induction medium). S: Scutellum, C: Coleorhiza, PR: Primary root.




Figure 4. Immature wheat embryo sections treated with germin DIG-labeled (A) anti-sense RNA probes and (B) sense RNA probes (after 6-hour incubation in callus induction medium). S: Scutellum, C: Coleorhiza, PR: Primary root.

formation during these 6-hour incubation period in callus induction medium; however, there was a shift in the quality of genes expressed in immature wheat embryos as the figures indicate (Figs. 3A and 4A).

The current study indicated that germin gene expression was induced during callus formation. *In situ* RNA hybridization, one of the most powerful techniques to characterize genes, revealed that germin genes were associated with the scutellum, coleorhiza and primary root during callus induction. In view of the current results, it seems that accumulation of germin genes during the early hours of callus induction might be related to the arrangement of the reorganization of cells for the transition from normal growth to callus formation in which cells retain their embryonic capacity. Furthermore, the results obtained in this study support Domon (19), who showed that germin accumulation was related to the cells which maintain their embryonic capacity during pine embryogenesis. Obviously, more studies are needed to establish the mechanisms involved in callus formation and embryonic capacity of cells.

References

1. Morris, P.C., Maddock, S.E., Jones, M.G.K. and Bowles, D.J., Changes in the levels of wheat- and barley-germ agglutinins during embryogenesis *in vivo*, *in vitro* and during germination. *Planta*, 166: 407-413, 1985.
2. Morris P.C., Kumar, A., Bowles, D.J. and Cuming, A.C., Osmotic stress and abscisic acid regulate expression of the wheat Em genes. *Eur. J. Biochem.*, 190: 625-630, 1990.
3. Quatrano, R.S., Ballo, B.L., Williamson, J.D., Hamblin, M.T. and Mansfield, M., In: *Plant Molecular Biology*, Goldberg, R. & Alan, R.L. (eds.), New York, 1983, 343-353.
4. Goldberg, R.B., Barker, S.J. and Perez-Grau, L., Regulation of gene expression during plant embryogenesis. *Cell*, 56: 149-160, 1989.
5. Bandurski, R.S., Schulze, A., Desrosiers, M., Jensa, P., Ebel, B. and Reinecke, D., In: *Plant Growth Substances*, Pharis, R.B. & Rood, S.B. (eds), Berlin, Heidelberg, New York, 1990 Springer-Verlag, 341-352.
6. R.S., In: *Oxford Surveys of Plant Molecular and Cell Biology*, Mifflin, B.J. ed., U K, 1986 Oxford University Press, 477.
7. Morris, P.C. and Bowles, D.J., In: *Growth Regulators and Seeds*, Monograph 15, U K, 1987 British Plant Growth Regulator Group, 65-76.
8. Fincher, G.B., Molecular and cellular biology associated with endosperm mobilization in germinating cereal grains. *Ann. Rev. Plant Physiol. Plant Mol. Biol.*, 40: 305-346, 1989.
9. Abeles, F.B., Morgan, P.W. and Saltveit, M.E. Jr., *Ethylene in plant biology*. 2nd edition, USA, 1992, Academic Press Inc.
10. Thompson, F.B. and Lane, B.G., Relation of protein synthesis in imbibing wheat embryos to the cell-free translational capacities of bulk mRNA from dry and imbibing embryos. *J. Biol. Chem.*, 255: 5965-5970, 1980.
11. Rahman, S., Grzelczak, Z.F., Kennedy, T.D. and Lane, B.G., Germin: molecular cloning of cDNA that selects germin mRNA from bulk wheat mRNA. *Biochem. Cell Biol.*, 66: 100-106, 1988.
12. Lane, B.G., Cellular desiccation and hydration: developmentally regulated proteins, and the maturation and germination of seed embryos. *FASEB J.*, 5: 2893-2901, 1991.
13. Hurkman, W.J., Tao, H.P., and Tanaka, C.K., Germin-like polypeptides increase in barley roots during salt stress. *Plant Physiol.*, 97: 366-374, 1991.
14. Hurkman, W.J. and Tanaka, C.K., Effect of salt stress on germin gene expression in barley roots. *Plant Physiol.*, 110: 971-977, 1996.
15. Dumas, B., Freyssinet, G. and Pallett, K.E., Tissue-specific expression of germin-like oxalate oxidase during development and fungal infection of barley seedlings. *Plant Physiol.*, 107: 1091-1096, 1995.
16. Zhang, Z., Collinge, D.B. and Thordal-Christensen, H., Germin-like oxalate oxidase, a H₂O₂-producing enzyme, accumulates in barley attacked by the powdery mildew fungus. *The Plant J.*, 8: 139-145, 1995.
17. Thordal-Christensen, H., Zhang, Z., Wei, Y. and Collinge, D.B., Subcellular localization of H₂O₂ in plants. H₂O₂ accumulation in papillae and hypersensitive response during the barley-powdery mildew interaction. *The Plant J.*, 11: 1187-1194, 1997.
18. Wei, Y., Zhang, Z., Andersen, C.H., Schmelzer, E., Gregersen, P.L., Collinge, D.B., Smedegaard-Petersen, V., and Thordal-Christensen, H., An epidermis/papilla-specific oxalate oxidase-like protein in the defence response of barley attacked by the powdery mildew fungus. *Plant Mol. Biol.*, 36: 101-112, 1998.

19. Domon, J.M., Dumas, B., Laine, E., Meyer, Y., David, A. and David, H., Three glycosylated polypeptides secreted by several embryogenic cell cultures of pine show highly specific serological affinity to antibodies directed against the wheat germin apoprotein monomer. *Plant Physiol.*, 108: 141-148, 1995.
20. Heintzen, C., Fischer, R., Melzer, S., Kappeler, S., Apel, K. and Staiger, D., Circadian oscillations of a transcript encoding a germin-like protein that is associated with cell walls in young leaves of the long-day plant *Sinapis alba* L. *Plant Physiol.*, 106: 905-915, 1994.
21. Ono, M., Sage-Ono, K., Inoue, M., Kamada, H. and Harada, H., Transient increase in the level of mRNA for a germin-like protein in leaves of the short-day plant *Pharbitis nil* during the photoperiodic induction of flowering. *Plant Cell Physiol.*, 37: 855-861, 1996.
22. Berna, A. and Bernier, F., Regulated expression of a wheat germin gene in tobacco: oxalate oxidase activity and apoplastic localization of the heterologous protein. *Plant Mol. Biol.*, 33: 1997.
23. Braun, H., Czihal, A., Shutov, A.D. and Baumlein, H., A vicilin-like seed protein of cycads: similarity to sucrose-binding proteins. *Plant Mol. Biol.*, 31: 35-44, 1996.
24. Dunwell, J.M., Cupins: A new superfamily of functionally diverse proteins that include germans and plant storage proteins. *Biotech. Genet. Enging. Rev.*, 15: 1-32, 1998.
25. Shutov, A.D., Braun, H., Chesnokov, Y.V. and Baumlein, H., A gene encoding a vicilin-like protein is specifically expressed in fern spores. Evolutionary pathway of seed storage globulins. *Eur. J. Biochem.*, 252: 79-89, 1998.
26. Berna, A. and Bernier, F., Regulation by biotic and abiotic stress of a wheat germin gene encoding oxalate oxidase, a H₂O₂-producing enzyme. *Plant Mol. Biol.*, 39: 539-549, 1999.
27. Dumas, B., Sailland, A., Cheviet, J-P., Freyssinet, G. and Pallett, K., Identification of barley oxalate oxidase as a germin-like protein. *C. R. Acad. Sci. Paris, Sciences de la vie/Life sciences*, 316: 793-798, 1993.
28. Lane, B.G., Dunwell, J.M., Ray, J.A., Schmitt, M.R., and Cuming, A.C., Germin, a protein marker of early plant development, is an oxalate oxidase. *J. Biol. Chem.*, 268: 12239-12242, 1993.
29. Luttrell, B.M., The biological relevance of the binding of calcium ions by inositol phosphates. *J. Biol. Chem.*, 268: 1521-1524, 1993.
30. Showalter, A.M., Structure and function of plant cell wall proteins. *The Plant Cell*, 5: 9-23, 1993.
31. Çalışkan, M., The metabolism of oxalic acid. *Tr. J. of Zoology*, 24: 103-106, 2000.
32. Çalışkan, M., Germin, which is an oxalate oxidase, takes function in many aspects of plant life. *Tr. J. of Biology*, 24: 717-724, 2000.
33. Çalışkan, M. and Cuming, A. C., Spatial specificity of H₂O₂ - generating oxalate oxidase gene expression during wheat embryo development. *Plant Journal*, 15: 165-171, 1998.
34. Çalışkan, M. and Cuming, A.C., Monitoring of germin synthesis by pre-absorbed anti-germin serum. *Tr. J. of Biology*, 24: 271-278, 2000.
35. Çalışkan, M. and Cuming, A.C., Temporal and spatial analysis of germin biosynthesis in wheat tissues. *Tr. J. of Biology*, 24: 775-782, 2000.
36. Maddock, S.E., Lancaster, V.A., Risiott, R. and Franklin, F., Plant regeneration from cultured immature embryos and inflorescences of 25 cultivars of wheat (*Triticum aestivum*). *J. Exp. Bot.*, 34: 915-926, 1983.
37. Vasil V., Castillo, A.M., Fromm, M.E. and Vasil, I.K., Herbicide resistant fertile transgenic wheat plants obtained by microprojectile bombardment of regenerable embryogenic callus. *Bio/Technology*, 10: 667-674, 1992.

38. Lane, B.G., Bernier, F., Dratewka-Kos, E., Shafai, R., Kennedy, T.D., Pyne, C., Munro, J.R., Vaughan, T., Walters, D., and Altomare, F., Homologies between members of the germin gene family in hexaploid wheat and similarities between these wheat germinals and certain *Physarum spherulins*. J. Biol. Chem., 266: 10461-10469, 1991.
39. Cox, K.H., Deleon, D.V., Angerer, L.M. and Angerer, R.C., Detection of mRNA in sea urchin embryos by *in situ* hybridization using asymmetric RNA probes. Developmental Biology, 101: 485-502, 1984.
40. Jackson, D.P., In: Molecular Plant Pathology: a practical approach, D.J. Bowles, S.J. Gurr, and M. McPherson (eds), U K, 1991 Oxford University Press.
41. Turet, M., Molecular analysis of changes in gene expression during callus induction from wheat embryos. Ph.D. Thesis, University of Leeds, 1993.